

LATE CRETACEOUS *PUPOIDES* PFEIFFER 1854 (GASTROPODA: PUPILLIDAE) FROM URUGUAY (QUEGUAY FORMATION)

F. CABRERA & S. MARTÍNEZ

Departamento de Paleontología, Facultad de Ciencias, Universidad de la República. Iguá 4225, P.C. 11400. Montevideo, Uruguay.

Abstract A new species of *Pupoides* Pfeiffer 1854, subgenus *Ischnopupoides* Pilsbry 1926, is described for the Late Cretaceous of Uruguay (Queguay Formation), being the oldest record of the genus and subgenus. *Pupoides* (I.) *gnocco* new species is characterized by a small dextral fusiform shell, constituted by a spire comprising five slightly convex whorls, oblicuous axial ornamentation, subrounded aperture, and an expanded outer lip that lacks dentition.

Key words Pupillidae, *Ischnopupoides*, Late Cretaceous, Uruguay

INTRODUCTION

The family Pupillidae (Turton 1831) comprises a clade of very small pulmonate snails distributed all over the world, mainly in Europe, North America and North Africa, but also recorded in Asia, Oceania and South America. *Pupoides* Pfeiffer 1854 is widely distributed throughout the world. In South America it is recognized in many regions in Recent and earlier periods (Hylton Scott, 1970; Stuardo & Vargas-Almonacid, 2000; Mächtle *et al.*, 2010). It is characterized by small pupoid to fusiform dextral shells with an oval to subrounded aperture, without lamellae in most species (Pfeiffer, 1854; Thiele, 1992); a few possess a small parietal tubercle (Schileyko, 1998).

The material herein described was found in the Queguay Formation (Late Cretaceous) of Uruguay (Fig. 1), which is composed by calcretes (pedogenic and groundwater), palustrine, and lacustrine limestones (Veroslavsky *et al.*, 1997; Martínez & Veroslavsky, 2004; Tófaló & Morrás, 2009; Alonso Zarza *et al.*, 2011). Fossils are found in the limestone beds that represent paleosols, that yield diverse terrestrial and freshwater gastropod assemblages. These assemblages include *Biomphalaria walteri*, *Eoborus charruanus*, *Bulimulus klappenbachii* and *Bahiensis priscus* (Martínez *et al.*, 1997; Martínez *et al.*, 2001; Cabrera & Martínez, 2012; Cabrera, 2015). Other fossils are characean oogons, ostracods, *Celtis santosi* (Ulmaceae) endocarps, ichnofossils such as *Celliforma* isp. and *Fictovichnus* isp. (hymenopteran nests and coleopteran pupal chambers respectively), and Neosauropoda

eggshells (Genise *et al.*, 2010; Alonso Zarza *et al.*, 2011; Verde, 2012; Cabrera, 2015).

The Holotype and most specimens are deposited in the Paleontological Collection of Facultad de Ciencias FCDP-I (Universidad de la República; Montevideo, Uruguay); other specimens are deposited in the Paleontological Collection of Centro de Estudios Aplicados del Litoral PZ-CTES (Universidad del Nordeste; Corrientes, Argentina).

SYSTEMATICS

Suprageneric classification follows Bouchet & Rocroi (2005).

Superfamily PUPILLOIDEA Turton 1831

Family PUPILLIDAE Turton 1831

Genus *Pupoides* Pfeiffer 1845

Type species *Bulimus nitidulus* Pfeiffer 1845

Subgenus *Ischnopupoides* Pilsbry 1926

Type species *Pupa hordacea* Gabb 1866

Pupoides (*Ischnopupoides*) *gnocco* sp. nov.

Pupoides sp.: Morton & Herbst, 1993

Vertigo sp.: Morton & Herbst, 1993

Pupillidae indet.: Martínez *et al.*, 2001

Holotype FCDP-I 7258 (Fig. 2A–C).

Type locality Uruguay, Canelones Department, Sauce Solo locality (34°24'03.4"S 55°38'37.7"W).

Paratype FCDP-I 7843 (Fig. 2D–F)

Paratype locality Uruguay, Paysandú Department, Quebracho locality (31°55'13.55"S 57°54'21.62"W)

Other material (n=40) FCDP – I 4662, 6471, 6475, 7242, 7262, 11 shells (Quebracho, Paysandú); 4665, 4666, 7225, 7264, 8 shells (Piedras Coloradas, Paysandú); 4679, 7219, 8 shells (Sauce Solo, Canelones); 7215, 3 shells (surroundings of Algorta Town, Río Negro); 7231, 1 shell (Palmar, Soriano); PZ-CTES 5340, 9 shells (Piedras Coloradas, Paysandú).

Holotype measurements (mm) height 3.57 (height range of 10 individuals: 2.8 to 4.15), width 1.84, last whorl 2.47; aperture height 1.70, width 1.32 (figure 2A–C).

Geographic and stratigraphic occurrence Sauce Solo (Canelones Department); surroundings of Algorta Town (Río Negro Department); Palmar (Soriano Department); Piedras Coloradas and Quebracho (Paysandú Department), Uruguay (Fig. 1). Queguay Formation, Late Cretaceous.



Figure 1 Map of Uruguay showing the locations where the species was found. 1 Quebracho, Paysandú; 2 Piedras Coloradas, Paysandú; 3 surroundings of Algorta Town, Río Negro; 4 Palmar, Soriano; 5 Sauce Solo, Canelones.

Etymology *gnocco* is the singular name for the unit of an Italian pasta (plural *gnocchi*). The shell shape of the new species resembles this traditional food.

Diagnosis Small dextral fusiform shell, with five slightly convex whorls; oblique axial ornamentation. Subrounded aperture, outer lip expanded without lamellae.

Description Shell small (near 4mm long) with a fusiform dextral shape, five slightly convex whorls; last whorl comprises 70% of the shell; shallow sutures; conspicuous slightly oblique and sigmoidal axial striations. Subrounded toothless aperture; thick and expanded outer lip.

Discussion The morphology of the studied specimens is consistent with the general diagnostic characters of Pupillidae Turton 1831. The particular features of our fossil material, either by their morphology and by their geographic occurrence, resemble those of other three genera included in the family: *Pupoides* Pfeiffer 1854, *Gastrocopta* Wollaston 1878 and *Pupilla* Fleming 1828. Species of Diplommatinidae (Pfeiffer 1857) were also considered, but rejected. Species of this family are small, some of them with a pupoid-fusiform shell, as in the new species. But the ornamentation is quite different, consisting of conspicuous separated ribs in Diplommatinidae, compared to narrow striae in the new species. Whorls and sutures are also different – in Diplommatinidae sutures are deep, and whorls are rounded and globose, while in the new species, whorls are less rounded, and sutures are shallow. In South America there is only one genus (with eight species) of this family: *Adelopoma* Doering, 1884, that have a sinistral shell (Hausdorf & Muñoz, 2004; Martins & Simone, 2014).

Pupilla is the most similar in shape to the new species, except for the absence of the characteristic parietal teeth in this genus. Recent work of Nekola *et al.* (2014) indicates that the aperture and the number of apertural teeth vary between individuals within species of *Pupilla*, so the lack of a parietal tooth in the new species would not be diagnostic. The new species differs from this genus by the ornamentation, and by the shape of whorls and sutures, along with the geographical distribution of this genus that does not currently include South America according to GBIF




Figure 2 A–C *Pupoides (Ischnopupoides) gnocco* sp. nov. Holotype, FCDP–I 7258. **A** apertural view, **B** lateral view, **C** posterior view. The aperture is reconstructed because it was broken while removing the carbonates that were encrusted on it. Photos and reconstruction by Gustavo Lecuona. **D–F** *Pupoides (Ischnopupoides) gnocco* sp. nov. Paratype, FCDP–I 7843. **D** apertural view, **E** lateral view, **F** posterior view. Scales: 1mm.

(<http://www.gbif.org/species/2294040>, visited on 01/15/15).

Gastrocopta is reported from many regions of South America, especially in Brazil and Uruguay (Scarabino, 2003; Simone, 2006). However, *Gastrocopta* has many teeth in the aperture and a folded outer lip (Wollaston, 1878; Roth, 1999), while the new species has no teeth, and the outer lip is smooth and expanded.

Pupoides has a wide geographic occurrence and, although many species have been described in the region (Argentina, Chile, Perú and Bolivia) (Hylton Scott, 1970; Stuardo & Vargas-Almonacid, 2000; Mächtle *et al.*, 2010), it had not been yet recorded in Uruguay. Its peculiar morphological characters are the presence of a fusiform to subconical shell, with five to six spire whorls ornamented with axial striae, and the development of an oval to subrounded aperture that can be toothless or have small teeth (Pfeiffer, 1854; Hylton Scott, 1970). In consequence, both the morphology and geographic proximity contribute to our placement of the new species within this genus. Moreover, they can be assigned to the subgenus *Ischnopupoides* Pilsbry 1926, which is recognized by the possession of small shells (less than 3.5mm) in most species, with a rounded aperture without teeth or with a small tubercle, and a spire devoid of sculpture or ornamented by thin striae parallel to the growth lines (Pilsbry, 1926; Schileyko, 1998; Benton *et al.*, 2015).

The type species of *Pupoides* is *Bulimus nitidulus* according to Pfeiffer (1845). This species has a small, turreted shell (5.1mm) of six whorls separated by deeply impressed sutures and an oval, toothless aperture. This species differs from *Pupoides (Ischnopupoides) gnocco* sp. nov. in the number of whorls (five in the new species), and the size and shape of the shell.

Pupa hordacea Gabb 1866 is the type species of *Ischnopupoides* (AMNH – http://research.amnh.org/iz/types_db/details.php?specimen_id=10296, visited on 07/05/2016), and taking into account the characters described originally, this species differs from *Pupoides (Ischnopupoides) gnocco* sp. nov. by the number of whorls (six in *P. hordacea*); the shell shape (being more elongated in *P. hordacea*) and because it is smaller than the fossil species, being almost 3mm in length (2.8mm).

Two extant species of *Pupoides (Ischnopupoides)* and two extant species of *Pupoides (Pupoides)* are recorded from Chile, Bolivia, Peru and Argentina

(Hylton Scott, 1970; Stuardo & Vargas-Almonacid, 2000). *Pupoides (Ischnopupoides) minimus* (Philippi 1860), from Chile, differs from the new species by the presence of a tubercle situated close to the top of the outer lip, and by the number of whorls (six whorls in *P. (I.) minimus*).

Pupoides (Ischnopupoides) paredesi (d'Orbigny 1835), inhabiting Chile, Peru, Bolivia and Argentina, is separated from the new species by its larger shell (5mm), with six whorls, and its large oval aperture.

Pupoides (Pupoides) chordatus (Pfeiffer 1857) from Argentina, may be distinguished from the new species by its almost conical – sub conical shell provided with an expanded last whorl, while the new Uruguayan taxon has a pupoid shell. *Pupoides (P.) centralis* (Hylton Scott 1970), also from Argentina, is similar to *P. (P.) chordatus*, therefore differs from the new species by the same characters.

Pupoides (P.) albilabris (Adams 1841) has been mentioned from the Pleistocene – Holocene of Perú (Mächtle *et al.*, 2010) and differs from *P. (I.) gnocco* sp. nov. by the shell-shape, which conical in *P. albilabris* and pupoid in the new species, and by the number of whorls, six in *P. albilabris* and five in *P. (I.) gnocco* sp. nov.

ACKNOWLEDGEMENTS

Lucía Vivanco found the complete specimen selected as Holotype. Verónica Espíndola allowed the access to the Paleontological Collection of the Centro de Estudios Aplicados del Litoral (Corrientes, Argentina). M. Verde, A. Batista, G. Roland, F. Montenegro and N. Batalla helped in field works. Gustavo Lecuona helped with pictures of specimens. J.C. Nekola provided bibliography and made valuable comments that helped to improve this paper. Claudia del Río helped with language. Project FCE 3_2011_1_5702 (ANII), PEDECIBA, and AUGM gave financial support. Finally, we wish to thank to the staff of the Paleontological Department that democratically voted for the name of the new species.

REFERENCES

- ADAMS CB 1841 *Catalogue of the land shells which inhabit Jamaica*.- Amherst, Mass. s.n. Pp. 2.
ALONSO-ZARZA AM, GENISE JF & VERDE M 2011 Sedimentology, diagenesis and ichnology of

- Cretaceous and Palaeogene calcretes and palustrine carbonates from Uruguay. *Sedimentary Geology* **236**: 45–61.
- AMNH – American Museum of Natural History 2002 – 2005. Accessed via http://research.amnh.org/iz/types_db/details.php?specimen_id=10296 on 2016-07-05
- BENTON RC, TERRY JR. DO, EVANOF E, McDONALD HG 2015 *The White River Badlands. Geology and Paleontology*. Indiana University Press, Indiana, United States.
- BOUCHET P & ROCROI JP 2005 Classification and Nomenclator of Gastropod Families. *Malacologia* **47**(1–2): 397.
- CABRERA F & MARTÍNEZ S 2012 The oldest Odontostomidae (Mollusca: Gastropoda): Bahiensis priscus n. sp. (Paleocene, Uruguay). *Paläontologische Zeitschrift* **86**: 451–456.
- CABRERA F 2015 Paleobiodiversidad y paleoecología de moluscos continentales. El Cretácico tardío – Paleógeno de Uruguay como caso de estudio. Master's Thesis in Biology. PEDECIBA Biología. Montevideo, Uruguay.
- FLEMING J 1828 *Exhibiting the descriptive characters and systematical arrangement of the genera and species of quadrupeds, birds, reptiles, fishes, mollusca, and radiata of the United Kingdom; including the indigenous, extirpated, and extinct kinds, together with periodical and occasional visitants*. Bell & Bradfute. Edinburgh, Scotland.
- GBIF SECRETARIAT: GBIF Backbone Taxonomy, 2013-07-01. Accessed via <http://www.gbif.org/species/2294040> on 2015-01-15
- GENISE JF, MELCHOR RN, BELLOSI ES & VERDE M 2010 Invertebrate and Vertebrate Trace Fossils from Continental Carbonates. *Developments in Sedimentology* **61**. Elsevier B.V.
- HAUSDORF B & MUÑOZ SG 2004 *Adelopoma peruvianum* new species from northern Peru (Gastropoda: Diplommatinidae). *Journal of Conchology* **38**(4): 369–372.
- HYLTON SCOTT MI 1970 El género Pupoides en Argentina (Pulmonata, Stylommatophora, Pupillidae). *Neotrópica* **16** (50): 89–90.
- MÄCHTLE B, UNKEL I, EITEL B, KROMER B & SCHIEGL S 2010 Molluscs as evidence for a late Pleistocene and early Holocene humid period in the southern coastal desert of Peru (14.5°S). *Quaternary Research* **73**: 39–47
- MARTÍNEZ S, VEROSLAVSKY G & VERDE M 1997 Primer registro del Paleoceno en el Uruguay: Paleosuelos calcáreos fosilíferos en la Cuenca de Santa Lucía. *Revista Brasileira de Geociências* **27**(3): 295–302.
- MARTÍNEZ S, VEROSLAVSKY G & VERDE M 2001 Paleoecología de los paleosuelos calcáreos fosilíferos («Calizas del Queguay» Paleoceno) de las regiones sur y litoral oeste del Uruguay. 11° Congreso Latinoamericano y 3er Uruguayo de Geología, Actas CD-Rom: Trabajo 219.
- MARTÍNEZ S & VEROSLAVSKY G 2004 Registros continentales del Terciario Temprano. In: Veroslavsky, G., Ubilla, M. & Martínez S. (Eds.): *Cuencas sedimentarias de Uruguay: geología, paleontología y recursos naturales – Cenozoico*: 63–82. – DIRAC-Facultad de Ciencias, Montevideo, Uruguay.
- MARTINS CM & SIMONE LRL 2014 A new species of *Adelopoma* from São Paulo Urban Park, Brazil (Caenogastropoda, Diplommatinidae) *Journal of Conchology* **41**(6): 765–773.
- MORTON LS & HERBST R 1993 Gastrópodos del Cretácico (Formación Mercedes) del Uruguay *Ameghiniana* **30**: 445–452.
- NEKOLA JC, COLES BF & HORSÁK M 2014 Species assignment in *Pupilla* (Gastropoda: Pulmonata: Pupillidae): integration of DNA-sequence data and conchology *Journal of Molluscan Studies* Advance Access 1–21. doi:10.1093/mollus/eyu083
- PFEIFFER L 1854 Zur Mollusken Fauna der Insel Cuba *Malakozoologischen Blätter* **1**: 192.
- PILSBRY, HA 1926 *Manual of Conchology. Pupillidae*. Second series, Pulmonata. Philadelphia **27**: 1–369.
- SCARABINO F 2003 Lista sistemática de los Gastrópoda terrestres vivientes de Uruguay. *Comunicaciones de la Sociedad Malacológica del Uruguay* **8**: 203–214.
- SCHILEYKO AA 1998 Treatise on Recent Terrestrial Pulmonate Molluscs. **1**: Achatinellidae, Amastridae, Orculidae, Strobilopsidae, Spelaeodiscidae, Valloniidae, Cochlicopidae, Pupillidae, Chondrinidae, Pyramidulidae. *Ruthenica Supplement 2*, Moscow, pp 112–113
- SIMONE LRL 2006 *Land and Freshwater Molluscs of Brazil*. Museu de Zoologia Universidade de São Paulo. São Paulo, Brazil.
- STUARDO JR & VARGAS-ALMONACID P 2000 Moluscos terrestres de Chile. Sinonimia y problemas relacionados: 1. Familias Veronicellidae, Pupillidae y Achatinellidae (GASTROPODA: PULMONATA). *Gayana* **64** (2) ISSN 07176538.
- ROTH B 1999 A new species of *Gastrocopta* (Gastropoda: Pulmonata: Pupillidae) from the Deep River Formation, Late Oligocene or Early Miocene, Montana. *The Veliger* **42**(3): 286–288.
- THIELE J 1992 Handbook of systematic malacology. Translation of: Thiele, J. (1929–1935) *Handbuch der systematischen Weichtierkunde*. Smithsonian Institution Libraries, Pauls Press, New Delhi, India.
- TÓFALO OR & MORRÁS HJM 2009 Evidencias paleoclimáticas en duricostras, paleosuelos y sedimentitas silicoclásticas, del Cenozoico de Uruguay *Revista de la Asociación Geológica Argentina* **65**(4): 674–686.
- TURTON W 1831 *A manual of the land and fresh-water shells of the British Islands, arranged according to the more modern systems of classification; and described from perfect specimens in the author's cabinet*. Printed by Richard Taylor, Red Lion Court, Fleet Street. London, England.
- VERDE M 2012 *ICnología de la Formación Asencio y las "Calizas Del Queguay"*. PhD thesis. PEDECIBA Biología, 204 pp. Universidad de la República, Montevideo.
- VEROSLAVSKY G, MARTÍNEZ S & DE SANTA ANA H 1997 Calcretas de aguas subterráneas y pedogénicas: génesis de los depósitos carbonáticos de la

Cuenca de Santa Lucía, sur del Uruguay (Cretácico Superior?-Paleógeno). *Revista de la Asociación Argentina de Sedimentología* 4(1): 25-35.

WOLLASTON TV 1878 *Testacea atlantica or the land and freshwater shells of the Azores, Madeiras, Salvages,*

Canaries, Cape Verdes, and Saint Helena. L. Reeve & Co. London, England.