



UNIVERSIDAD
DE LA REPÚBLICA
URUGUAY



FACULTAD DE
INGENIERÍA
UDELAR



Programa de
Desarrollo de las
Ciencias Básicas

Coding of Multichannel Signals with Irregular Sampling Rates and Data Gaps

Pablo Cerveñansky Fierro

Tesis de Maestría presentada a la Facultad de Ingeniería de la Universidad de la República
en cumplimiento parcial de los requerimientos para la obtención del título de
Magister en Informática

Tutores

Dr. Álvaro Martín
Dr. Gadiel Seroussi

Tribunal

Dra. Lorena Etcheverry
Dr. Jorge Silva
Dr. Leonardo Steinfeld

Montevideo, Uruguay
Julio de 2021

Introduction

This document includes each plot figure generated from the experimental results obtained in our work. In **Appendix A** we present the figures corresponding to Section 4.2 (Comparison of Masking and Non-Masking Variants). In **Appendix B** we present the figures corresponding to Section 4.3 (Window Size Parameter). Finally, in **Appendix C** we present the figures corresponding to Section 4.4 (Algorithm Compression Performance).

Table of Contents

Introduction	ii
Table of Contents	ii
List of Figures	iv
A Figures: Comparison of Masking and Non-Masking Variants	1
B Figures: Window Size Parameter	23
C Figures: Algorithm Compression Performance	49

List of Figures

A.1	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “VWC” of the dataset IRKIS.	2
A.2	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “SST” of the dataset SST. In the RD plot for algorithm PCA we highlight with a red circle the marker for the maximum value (50.78%) obtained for all the tested CAIs.	3
A.3	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Velocity” of the dataset ADCP.	4
A.4	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “GHI” of the dataset Solar.	5
A.5	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “DNI” of the dataset Solar.	6
A.6	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “DHI” of the dataset Solar.	7
A.7	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset ElNino.	8
A.8	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset ElNino.	9
A.9	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Zon. Wind” of the dataset ElNino.	10
A.10	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Mer. Wind” of the dataset ElNino.	11
A.11	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Humidity” of the dataset ElNino.	12
A.12	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Air Temp.” of the dataset ElNino.	13
A.13	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Sea Temp.” of the dataset ElNino.	14
A.14	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Hail.	15
A.15	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Hail.	16
A.16	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Size” of the dataset Hail.	17
A.17	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Tornado.	18
A.18	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Tornado. In the RD plot for algorithm APCA we highlight with a blue circle the marker for the minimum value (-0.29%) obtained for all the tested CAIs.	19
A.19	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Wind.	20
A.20	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Wind.	21

A.21	CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Speed” of the dataset Wind.	22
B.1	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1202.csv” of the dataset IRKIS.	24
B.2	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1203.csv” of the dataset IRKIS. In the RD plot for variant PCA_M we highlight with a red circle the marker for the maximum value (10.6%) obtained for all the tested CAIs.	25
B.3	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1204.csv” of the dataset IRKIS.	26
B.4	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1205.csv” of the dataset IRKIS.	27
B.5	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-222.csv” of the dataset IRKIS.	28
B.6	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-333.csv” of the dataset IRKIS.	29
B.7	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-SLF2.csv” of the dataset IRKIS.	30
B.8	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-01-2017.csv” of the dataset SST.	31
B.9	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-02-2017.csv” of the dataset SST.	32
B.10	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-03-2017.csv” of the dataset SST.	33
B.11	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-01-2015.csv” of the dataset SST.	34
B.12	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-02-2015.csv” of the dataset SST.	35
B.13	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-03-2015.csv” of the dataset SST.	36
B.14	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2011.csv” of the dataset Solar.	37
B.15	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2011.csv” of the dataset Solar.	38
B.16	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2011.csv” of the dataset Solar.	39

B.17	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2012.csv” of the dataset Solar.	40
B.18	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2012.csv” of the dataset Solar.	41
B.19	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2012.csv” of the dataset Solar.	42
B.20	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2013.csv” of the dataset Solar.	43
B.21	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2013.csv” of the dataset Solar.	44
B.22	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2013.csv” of the dataset Solar.	45
B.23	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2014.csv” of the dataset Solar.	46
B.24	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2014.csv” of the dataset Solar.	47
B.25	Plots of w_{global}^* , w_{local}^* , and the RD between $c_{\langle a_v, w_{global}^*, e \rangle}$ and $c_{\langle a_v, w_{local}^*, e \rangle}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2014.csv” of the dataset Solar.	48
C.1	CR and window size parameter plots for every evaluated algorithm, for the data type “VWC” of the dataset IRKIS.	50
C.2	CR and window size parameter plots for every evaluated algorithm, for the data type “SST” of the dataset SST.	51
C.3	CR and window size parameter plots for every evaluated algorithm, for the data type “Velocity” of the dataset ADCP.	52
C.4	CR and window size parameter plots for every evaluated algorithm, for the data type “GHI” of the dataset Solar.	53
C.5	CR and window size parameter plots for every evaluated algorithm, for the data type “DNI” of the dataset Solar.	54
C.6	CR and window size parameter plots for every evaluated algorithm, for the data type “DHI” of the dataset Solar.	55
C.7	CR and window size parameter plots for every evaluated algorithm, for the data type “Latitude” of the dataset ElNino.	56
C.8	CR and window size parameter plots for every evaluated algorithm, for the data type “Longitude” of the dataset ElNino.	57
C.9	CR and window size parameter plots for every evaluated algorithm, for the data type “Zon. Wind” of the dataset ElNino.	58
C.10	CR and window size parameter plots for every evaluated algorithm, for the data type “Mer. Wind” of the dataset ElNino.	59
C.11	CR and window size parameter plots for every evaluated algorithm, for the data type “Humidity” of the dataset ElNino.	60
C.12	CR and window size parameter plots for every evaluated algorithm, for the data type “Air Temp.” of the dataset ElNino.	61

C.13 CR and window size parameter plots for every evaluated algorithm, for the data type “Sea Temp.” of the dataset ElNino. For each error parameter $e \in E$, we use blue circles to highlight the markers for the minimum CR value and the best window size (in the respective plots corresponding to the best coding variant).	62
C.14 CR and window size parameter plots for every evaluated algorithm, for the data type “Latitude” of the dataset Hail.	63
C.15 CR and window size parameter plots for every evaluated algorithm, for the data type “Longitude” of the dataset Hail.	64
C.16 CR and window size parameter plots for every evaluated algorithm, for the data type “Size” of the dataset Hail.	65
C.17 CR and window size parameter plots for every evaluated algorithm, for the data type “Latitude” of the dataset Tornado.	66
C.18 CR and window size parameter plots for every evaluated algorithm, for the data type “Longitude” of the dataset Tornado.	67
C.19 CR and window size parameter plots for every evaluated algorithm, for the data type “Latitude” of the dataset Wind.	68
C.20 CR and window size parameter plots for every evaluated algorithm, for the data type “Longitude” of the dataset Wind.	69
C.21 CR and window size parameter plots for every evaluated algorithm, for the data type “Speed” of the dataset Wind.	70

Appendix A

Figures: Comparison of Masking and Non-Masking Variants




FIGURE A.1: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “VWC” of the dataset IRKIS.




FIGURE A.2: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type "SST" of the dataset SST. In the RD plot for algorithm PCA we highlight with a red circle the marker for the maximum value (50.78%) obtained for all the tested CAIs.




FIGURE A.3: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type "Velocity" of the dataset ADCP.




FIGURE A.4: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type "GHI" of the dataset Solar.




FIGURE A.5: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “DNI” of the dataset Solar.




FIGURE A.6: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “DHI” of the dataset Solar.




FIGURE A.7: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset ElNino.




FIGURE A.8: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset El Niño.




FIGURE A.9: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Zon. Wind” of the dataset El Niño.




FIGURE A.10: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Mer. Wind” of the dataset ElNino.




FIGURE A.11: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Humidity” of the dataset ElNino.




FIGURE A.12: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type "Air Temp." of the dataset ElNino.




FIGURE A.13: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type "Sea Temp." of the dataset ElNino.




FIGURE A.14: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Hail.




FIGURE A.15: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Hail.




FIGURE A.16: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Size” of the dataset Hail.




FIGURE A.17: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Tornado.




FIGURE A.18: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Tornado. In the RD plot for algorithm APCA we highlight with a blue circle the marker for the minimum value (-0.29%) obtained for all the tested CAIs.




FIGURE A.19: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Latitude” of the dataset Wind.




FIGURE A.20: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Longitude” of the dataset Wind.




FIGURE A.21: CR and RD plots for variants a_M and a_{NM} , for each algorithm $a \in A_M$, for the data type “Speed” of the dataset Wind.

Appendix B

Figures: Window Size Parameter




FIGURE B.1: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1202.csv” of the dataset IRKIS.




FIGURE B.2: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1203.csv” of the dataset IRKIS. In the RD plot for variant PCA_M we highlight with a red circle the marker for the maximum value (10.6%) obtained for all the tested CAIs.




FIGURE B.3: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1204.csv” of the dataset IRKIS.




FIGURE B.4: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-1205.csv” of the dataset IRKIS.




FIGURE B.5: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-222.csv” of the dataset IRKIS.




FIGURE B.6: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-333.csv” of the dataset IRKIS.




FIGURE B.7: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “VWC” of the file “irkis-SLF2.csv” of the dataset IRKIS.




FIGURE B.8: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-01-2017.csv” of the dataset SST.




FIGURE B.9: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-02-2017.csv” of the dataset SST.




FIGURE B.10: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “SST” of the file “sst-03-2017.csv” of the dataset SST.




FIGURE B.11: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-01-2015.csv” of the dataset SST.




FIGURE B.12: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-02-2015.csv” of the dataset SST.




FIGURE B.13: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “Velocity” of the file “adcp-03-2015.csv” of the dataset SST.




FIGURE B.14: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2011.csv” of the dataset Solar.




FIGURE B.15: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2011.csv” of the dataset Solar.




FIGURE B.16: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2011.csv” of the dataset Solar.




FIGURE B.17: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2012.csv” of the dataset Solar.




FIGURE B.18: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2012.csv” of the dataset Solar.




FIGURE B.19: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2012.csv” of the dataset Solar.




FIGURE B.20: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2013.csv” of the dataset Solar.




FIGURE B.21: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DNI” of the file “solar-2013.csv” of the dataset Solar.




FIGURE B.22: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2013.csv” of the dataset Solar.




FIGURE B.23: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “GHI” of the file “solar-2014.csv” of the dataset Solar.




FIGURE B.24: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type "DNI" of the file "solar-2014.csv" of the dataset Solar.




FIGURE B.25: Plots of w_{global}^* , w_{local}^* , and the RD between $c_{<a_v, w_{global}^*, e>}$ and $c_{<a_v, w_{local}^*, e>}$, as a function of the error parameter e , obtained for the data type “DHI” of the file “solar-2014.csv” of the dataset Solar.

Appendix C

Figures: Algorithm Compression Performance




FIGURE C.1: CR and window size parameter plots for every evaluated algorithm, for the data type “VWC” of the dataset IRKIS.




FIGURE C.2: CR and window size parameter plots for every evaluated algorithm, for the data type "SST" of the dataset SST.







FIGURE C.3: CR and window size parameter plots for every evaluated algorithm, for the data type “Velocity” of the dataset ADCP.










FIGURE C.7: CR and window size parameter plots for every evaluated algorithm, for the data type "Latitude" of the dataset ElNino.




FIGURE C.8: CR and window size parameter plots for every evaluated algorithm, for the data type "Longitude" of the dataset ElNino.




FIGURE C.9: CR and window size parameter plots for every evaluated algorithm, for the data type “Zon. Wind” of the dataset El Niño.




FIGURE C.10: CR and window size parameter plots for every evaluated algorithm, for the data type "Mer. Wind" of the dataset ElNino.




FIGURE C.11: CR and window size parameter plots for every evaluated algorithm, for the data type "Humidity" of the dataset ElNino.




FIGURE C.12: CR and window size parameter plots for every evaluated algorithm, for the data type "Air Temp." of the dataset ElNino.




FIGURE C.13: CR and window size parameter plots for every evaluated algorithm, for the data type “Sea Temp.” of the dataset ElNino. For each error parameter $e \in E$, we use blue circles to highlight the markers for the minimum CR value and the best window size (in the respective plots corresponding to the best coding variant).





FIGURE C.14: CR and window size parameter plots for every evaluated algorithm, for the data type "Latitude" of the dataset Hail.






FIGURE C.16: CR and window size parameter plots for every evaluated algorithm, for the data type “Size” of the dataset Hail.




FIGURE C.17: CR and window size parameter plots for every evaluated algorithm, for the data type “Latitude” of the dataset Tornado.




FIGURE C.18: CR and window size parameter plots for every evaluated algorithm, for the data type "Longitude" of the dataset Tornado.





FIGURE C.19: CR and window size parameter plots for every evaluated algorithm, for the data type "Latitude" of the dataset Wind.






FIGURE C.21: CR and window size parameter plots for every evaluated algorithm, for the data type “Speed” of the dataset Wind.