



# The permeability of human red blood cell membranes to hydrogen peroxide is independent of aquaporins

Received for publication, August 29, 2021, and in revised form, December 10, 2021 Published, Papers in Press, December 18, 2021, <https://doi.org/10.1016/j.jbc.2021.101503>

Florencia Orrico<sup>1,2,3,†</sup>, Ana C. Lopez<sup>1,2,3,†</sup>, Daniela Saliwonzky<sup>2,4</sup>, Cecilia Acosta<sup>2,4</sup>, Ismael Rodriguez-Grecco<sup>4</sup>, Isabelle Mouro-Chanteloup<sup>5,6</sup>, Mariano A. Ostuni<sup>5,6</sup>, Ana Denicola<sup>1,3</sup>, Leonor Thomson<sup>2,3,\*</sup>, and Matias N. Möller<sup>1,3,\*</sup>

From the <sup>1</sup>Laboratorio de Físicoquímica Biológica, Instituto de Química Biológica, Facultad de Ciencias, <sup>2</sup>Laboratorio de Enzimología, Instituto de Química Biológica, Facultad de Ciencias, <sup>3</sup>Centro de Investigaciones Biomédicas (CEINBIO), and <sup>4</sup>Departamento de Medicina Transfusional, Hospital de Clínicas, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay; <sup>5</sup>UMR\_S1134, BIGR, Inserm, Université de Paris, Paris, France; <sup>6</sup>Laboratoire d'Excellence GR-Ex, Paris, France

Edited by Dennis Voelker

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) not only is an oxidant but also is an important signaling molecule in vascular biology, mediating several physiological functions. Red blood cells (RBCs) have been proposed to be the primary sink of H<sub>2</sub>O<sub>2</sub> in the vasculature because they are the main cellular component of blood with a robust antioxidant defense and a high membrane permeability. However, the exact permeability of human RBC to H<sub>2</sub>O<sub>2</sub> is neither known nor is it known if the mechanism of permeation involves the lipid fraction or protein channels. To gain insight into the permeability process, we measured the partition constant of H<sub>2</sub>O<sub>2</sub> between water and octanol or hexadecane using a novel double-partition method. Our results indicated that there is a large thermodynamic barrier to H<sub>2</sub>O<sub>2</sub> permeation. The permeability coefficient of H<sub>2</sub>O<sub>2</sub> through phospholipid membranes containing cholesterol with saturated or unsaturated acyl chains was determined to be  $4 \times 10^{-4}$  and  $5 \times 10^{-3}$  cm s<sup>-1</sup>, respectively, at 37 °C. The permeability coefficient of human RBC membranes to H<sub>2</sub>O<sub>2</sub> at 37 °C, on the other hand, was  $1.6 \times 10^{-3}$  cm s<sup>-1</sup>. Different aquaporin-1 and aquaporin-3 inhibitors proved to have no effect on the permeation of H<sub>2</sub>O<sub>2</sub>. Moreover, human RBCs devoid of either aquaporin-1 or aquaporin-3 were equally permeable to H<sub>2</sub>O<sub>2</sub> as normal human RBCs. Therefore, these results indicate that H<sub>2</sub>O<sub>2</sub> does not diffuse into RBCs through aquaporins but rather through the lipid fraction or a still unidentified membrane protein.

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) is probably one of the most abundant reactive species derived from oxygen in biology. It can be produced enzymatically by several oxidases and also as a byproduct of mitochondrial respiration (1). Cells are equipped to deal with excess H<sub>2</sub>O<sub>2</sub> with several enzymatic antioxidant defenses, including peroxiredoxins, glutathione peroxidases, and catalases. H<sub>2</sub>O<sub>2</sub>-triggered signaling has been associated to various physiological responses, such as cell

migration, growth, and proliferation (1). It has been proposed that the signaling is mediated by a redox relay system, involving the oxidation of a peroxiredoxin that then conveys the oxidation equivalents to a second protein, likely with the assistance of an adapter protein (2, 3).

In the vasculature, H<sub>2</sub>O<sub>2</sub> is produced mainly by NADPH oxidases from endothelial cells. Once formed, it can diffuse and cause vasodilation (4, 5) or promote cell growth, proliferation, and migration in endothelial and smooth muscle cells (6, 7). There have been several attempts to measure the steady-state concentration of H<sub>2</sub>O<sub>2</sub> in blood. A normal range of 1 to 5 μM, which could increase up to 50 μM under inflammatory conditions, has been estimated (8). However, the great variations in the reported values indicate that it is a challenging task. Very likely, the actual concentration of H<sub>2</sub>O<sub>2</sub> is much lower because red blood cells (RBCs) consume H<sub>2</sub>O<sub>2</sub> rapidly and efficiently. In fact, RBCs contain a robust antioxidant defense consisting of peroxiredoxin 2 (Prx2), glutathione peroxidase 1, and catalase (9). Physiological low amounts of H<sub>2</sub>O<sub>2</sub>, including those produced by oxyhemoglobin (HbO<sub>2</sub>) autoxidation, are consumed mainly by Prx2, which is very abundant (240–520 μM) (10, 11) and reacts very rapidly ( $k = 1 \times 10^8$  M<sup>-1</sup>s<sup>-1</sup>, (12)). The reduction of Prx2 back to the active state by thioredoxin and thioredoxin reductase is limited by the low amount of NADPH present in the RBC, so large amounts of H<sub>2</sub>O<sub>2</sub> can transiently inactivate Prx2 (9, 13). In such a scenario, catalase consumes most of the remaining H<sub>2</sub>O<sub>2</sub>, at a slower rate (9).

H<sub>2</sub>O<sub>2</sub> is consumed by enzymes that are located in the cytosol of RBCs; therefore, it can cross the plasma membrane. Notably, the permeability of human RBC membrane to H<sub>2</sub>O<sub>2</sub> has not been determined but has been estimated in a few studies to be  $7 \times 10^{-4}$  and  $1.1 \times 10^{-3}$  cm s<sup>-1</sup> (9, 14), respectively. The permeability of RBCs from rat and horse has been determined to be  $1.2 \times 10^{-2}$  cm s<sup>-1</sup> and  $6 \times 10^{-4}$  cm s<sup>-1</sup>, respectively (15, 16), indicating a significant variability and prompting the determination of the permeability of human RBCs to H<sub>2</sub>O<sub>2</sub>. The permeability of several cells to H<sub>2</sub>O<sub>2</sub> has been quantitatively determined and found to range from

<sup>†</sup> These authors contributed equally to this work.

\* For correspondence: Matias N. Möller, [mmoller@fcien.edu.uy](mailto:mmoller@fcien.edu.uy); Leonor Thomson, [lthomson@fcien.edu.uy](mailto:lthomson@fcien.edu.uy).

## Permeability of membranes to hydrogen peroxide

$2 \times 10^{-4} \text{ cm s}^{-1}$  in Jurkat T cells to  $1.6 \times 10^{-3} \text{ cm s}^{-1}$  in human umbilical vein endothelial cells (17).

The role of aquaporins (AQPs) in facilitating the transport of  $\text{H}_2\text{O}_2$  has been studied qualitatively in different systems. It was found that not all AQPs, but certain isoforms later termed peroxiporins, facilitate the transport of  $\text{H}_2\text{O}_2$  (18). Among these are human aquaporin 3 (AQP3) and human aquaporin 8 (AQP8) (18–22). Human aquaporin 1 (AQP1), on the contrary, did not appear to facilitate  $\text{H}_2\text{O}_2$  transport (18, 22), but this has recently been disputed and remains controversial (23, 24). AQP1 is one of the most abundant proteins in the membrane of RBCs, and AQP3 is also present at lower levels (11), so  $\text{H}_2\text{O}_2$  may diffuse across the RBC membrane through these channels.

In contrast to the information on permeability of cellular membranes to  $\text{H}_2\text{O}_2$ , little is known about the permeability of lipid-only membranes that would mimic the lipid fraction of cell membranes. In these membranes, permeation would occur by simple diffusion, a process known to be greatly affected by solubility of the permeant molecule in the lipid phase (25, 26). Some reports have shown changes in diffusion rates depending on liposome composition, temperature, and compressibility, but no permeability coefficient ( $P_m$ ) values have been determined (15, 27, 28). Molecular dynamics simulations have provided detailed information on  $\text{H}_2\text{O}_2$  distribution across the lipid bilayer, but with no experimental counterpart to validate it (29–31). This lack of information somehow leads to the idea that  $\text{H}_2\text{O}_2$  can only traverse cell membranes by facilitated transport through AQPs, disregarding simple diffusion.

Herein, we first determined the solubility of  $\text{H}_2\text{O}_2$  in organic solvents representing different depths in the membrane, to gain insight of the thermodynamic barrier to  $\text{H}_2\text{O}_2$  diffusion across lipid membranes. To measure the  $P_m$  of lipid membranes, we used liposomes made of cholesterol (Chol) and phospholipids containing saturated or unsaturated acyl chains, which encapsulated catalase. The  $P_m$  of these membranes to  $\text{H}_2\text{O}_2$  was determined by comparing the rate of decomposition of  $\text{H}_2\text{O}_2$  by intact *versus* disrupted liposomes, using the enzyme latency principle. The same approach was used to determine the  $P_m$  of human RBCs to  $\text{H}_2\text{O}_2$ . Finally, we studied the permeation mechanism of  $\text{H}_2\text{O}_2$  in human RBCs, evaluating in the process the potential role of AQPs in  $\text{H}_2\text{O}_2$  transport.

## Results

### Solubility of $\text{H}_2\text{O}_2$ in organic solvents

As a first approximation to the solubility of  $\text{H}_2\text{O}_2$  in the lipid membrane, we used organic solvents that have been previously used to mimic several physicochemical properties of lipid membranes, namely *n*-octanol and hexadecane (25, 32, 33). As

expected, the solubility of  $\text{H}_2\text{O}_2$  in organic solvents was significantly lower than in water. The solubility in *n*-octanol was 15 times lower than in water, whereas in hexadecane, 122,000 times lower than in water (Table 1). Temperature-dependent assays further indicate that there is a greater contribution of enthalpy than entropy to the transfer of  $\text{H}_2\text{O}_2$  from water to these solvents, consistent with an important loss of hydrogen bonds of  $\text{H}_2\text{O}_2$  on transferring from water to the organic phase (Table 1).

Hexadecane and *n*-octanol differ in polarity as evidenced by the different dielectric constants (Table 1) and can be considered to represent different regions of the membrane (34–36). The relatively more polar *n*-octanol is similar to the acyl region near the carbonyl groups, whereas the less polar hexadecane is similar to the middle-bilayer region, thus a solubility profile across the membrane can be estimated. Figure 1 shows the estimated solubility profile across the membrane as well as the Gibbs energy of partition ( $\Delta G^\circ$ ) associated to such position. The profile is discontinuous between the membrane and bulk water because we have no experimental estimation on the solubility of  $\text{H}_2\text{O}_2$  in the highly polar headgroup region or the structured water region near the headgroups.

The  $P_m$  can be calculated in an inhomogeneous media such as the membrane as the inverse of the sum of resistances to permeation at the different depths through the membrane (37, 38):

$$P_m = \left( \int_{-d/2}^{d/2} \frac{1}{K(z)D(z)} dz \right)^{-1} \quad (1)$$

where  $K(z)$  and  $D(z)$  are the  $K_D^\circ$  and diffusion coefficient at a depth  $z$  in the membrane. To estimate  $P_m$ , it was assumed that  $D$  for  $\text{H}_2\text{O}_2$  is the same across all the membrane and equal to that in water ( $D = 1.4 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ , (39)). Two regions with different solubility were considered, the central region, 1.6 nm wide, where  $K_D^\circ$  is given by hexadecane, and two equal regions on either side, 0.7 nm wide each, where  $K_D^\circ$  is given by octanol. The headgroup region or the structured water region adjacent to the headgroups was not considered. The estimated  $P_m$  is  $7 \times 10^{-4} \text{ cm s}^{-1}$ . Of course, this is a simplistic estimation that does not take into account properties of lipid bilayers such as lipid packing and composition, and so experimental determinations of  $P_m$  were done next.


### Permeability of lipid membranes to $\text{H}_2\text{O}_2$

The permeability to  $\text{H}_2\text{O}_2$  was first measured in membranes containing dimiristoylphosphatidylcholine (DMPC) (0.4 mole

**Table 1**  
Partition constants of  $\text{H}_2\text{O}_2$  between organic solvents and water at 25 °C

Organic solvent	Dielectric constant	$K_D^\circ$	$\Delta G^\circ$ (kJ mol <sup>-1</sup> )	$\Delta H^\circ$ (kJ mol <sup>-1</sup> )	$\Delta S^\circ$ (J mol <sup>-1</sup> K <sup>-1</sup> )
Octanol	10.3	$(6.6 \pm 0.4) \times 10^{-2}$	6.7	5.9	-2.6
Hexadecane	2.05	$(8.2 \pm 0.6) \times 10^{-6}$	29.0	26.4	-8.7

For comparison, the dielectric constant of water is 78.4.



**Figure 1.** Estimated solubility profile of  $\text{H}_2\text{O}_2$  across the lipid bilayer, according to  $K_D^0$  (blue) and  $\Delta G^0$  (red). Assuming a uniform diffusion coefficient of  $\text{H}_2\text{O}_2$  across the membrane ( $D = 1.4 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$ , (39)), and using Equation 1, a  $P_m = 7 \times 10^{-4} \text{ cm s}^{-1}$  is estimated.  $\text{H}_2\text{O}_2$ , hydrogen peroxide.


fraction), as the base phospholipid, dipalmitoylphosphatidylglycerol (DPPG) (0.1 mole fraction), to provide a net negative charge to the vesicles and prevent them from fusing (40), and Chol (0.5 mole fraction), to provide mechanical stability and mimic typical mammal membrane composition. It was found that catalase encapsulated in liposomes caused  $\text{H}_2\text{O}_2$  decomposition in a dose-dependent manner, where the slope is the pseudo-first-order rate constant  $k_{\text{lipo}}$  (Fig. 2). The disruption of these liposomes by repeated extrusion through 30 nm pore filters resulted in an increase in the rate of  $\text{H}_2\text{O}_2$  decomposition, consistent with release of encapsulated catalase, re-equilibration between external and internal volumes, and the loss of the permeability barrier. The rate of  $\text{H}_2\text{O}_2$  decomposition was also dose dependent and used to calculate  $k_{\text{dis}}$  (Fig. 2). The ratio between  $k_{\text{lipo}}$  and  $k_{\text{dis}}$  yields  $R_{\text{H}_2\text{O}_2}$ .

The  $P_m$  was determined based on the enzyme latency method (16, 41, 42). Briefly, it is calculated considering that at steady state, the rate of  $\text{H}_2\text{O}_2$  diffusion through the membrane into the liposome will be equal to the sum of the rates of  $\text{H}_2\text{O}_2$

diffusion out of the liposomes and consumption by catalase inside the liposome (41). The advantage of the latency method is that the rate of the enzymatic reaction does not have to be much higher than the diffusion rate across the membrane but only slightly higher or in the order. It is considered that  $\text{H}_2\text{O}_2$  can diffuse into the vesicle and react with catalase and part of it can diffuse back to the external solution. A steady state is achieved rapidly, and then a competition between catalase decomposition and diffusion ensues, which generates a concentration gradient of  $\text{H}_2\text{O}_2$  across the membrane. The gradient of concentration of  $\text{H}_2\text{O}_2$  formed across the membrane is the inverse of  $R_{\text{H}_2\text{O}_2}$ .

A detailed explanation of the enzyme latency method is given in the Supporting Information. The final equation used to calculate  $P_m$  is given in Equation 2.

$$P_m = \frac{k_{\text{catalase}} R_{\text{H}_2\text{O}_2}}{\frac{A}{V}(1 - R_{\text{H}_2\text{O}_2})} \quad (2)$$



**Figure 2.** Permeability of  $\text{H}_2\text{O}_2$  through DMPC:DPPG:Chol (4:1:5) membranes. *A*, initial rate determination of 10 mM  $\text{H}_2\text{O}_2$  consumption by liposome-encapsulated catalase performed with increasing volumes of liposomes (5–25  $\mu\text{l}$ , red tones) and released catalase in disrupted liposomes (5–25  $\mu\text{l}$ , blue tones) at 25 °C. *B*, the initial rates are used to obtain pseudo-first-order constants for  $\text{H}_2\text{O}_2$  consumption,  $k_{\text{lipo}}$  for intact (red circles) and  $k_{\text{dis}}$  for disrupted (blue squares) liposomes. The ratio  $k_{\text{lipo}}/k_{\text{dis}}$  yields  $R_{\text{H}_2\text{O}_2}$ , which is used to calculate  $P_m$  according to Equation 2. *C*, activation energy of  $\text{H}_2\text{O}_2$  permeation determined in the 20 to 40 °C range,  $E_a = 57 \pm 8 \text{ kJ mol}^{-1}$  ( $n = 3$ ). Chol, cholesterol; DMPC, dimiristoylphosphatidylcholine; DPPG, dipalmitoylphosphatidylglycerol;  $\text{H}_2\text{O}_2$ , hydrogen peroxide.

## Permeability of membranes to hydrogen peroxide

where  $A/V$  is the surface area to internal volume ratio, calculated from the hydrodynamic radii ( $A/V \approx 3/r$ , disregarding the contribution of water to the hydrodynamic radius of these liposomes, calculated to be less than 4%). The rate constant  $k_{\text{catalase}}$  is the pseudo-first-order rate constant of catalase inside the liposome and was determined from the catalase work solution used to prepare the liposomes. It was assumed that the concentration of catalase inside the liposomes was the same as in the catalase work solution. It was a reasonable assumption because the liposomes were prepared and used the same day, and they were found to be stable (same catalase activity) for at least 3 days. Furthermore, the resulting  $P_m$  values were found to be consistent between batches from different preparations.

At 25 °C, it was found that  $P_m = (3.7 \pm 0.5) \times 10^{-5} \text{ cm s}^{-1}$  for liposomes made of DMPC, DPPG, and Chol (DMPC:DPPG:Chol, 4:1:5 mole fraction), whereas this value increased to  $(1.3 \pm 0.7) \times 10^{-3} \text{ cm s}^{-1}$  for liposomes made of dioleoylphosphatidylcholine (DOPC), 1-palmitoyl-2-oleoyl-phosphatidylglycerol, and Chol (DOPC:POPG:Chol, 4:1:5 mole fraction), pointing that the permeability to  $\text{H}_2\text{O}_2$  depended on the degree of unsaturation of the membrane lipids. The permeability increased with temperature. At 37 °C,  $P_m = (4.1 \pm 0.5) \times 10^{-4}$  and  $(5.5 \pm 0.3) \times 10^{-3} \text{ cm s}^{-1}$  for DMPC:DPPG:Chol (4:1:5) and DOPC:POPG:Chol (4:1:5), respectively (Table 2). The Arrhenius activation energy ( $E_a$ ) of the permeation process was calculated for each membrane and found to be  $130 \pm 20$  and  $57 \pm 8 \text{ kJ mol}^{-1}$  for DMPC:DPPG:Chol (4:1:5) and DOPC:POPG:Chol (4:1:5), respectively (Table 2).

The permeability of membranes to  $\text{H}_2\text{O}_2$  greatly depended on lipid composition. Membranes composed of DOPC:POPG:Chol (4:1:5) were 1 to 2 orders of magnitude more permeable to  $\text{H}_2\text{O}_2$  than membranes composed of DMPC:DPPG:Chol (4:1:5), suggesting that unsaturations in the acyl chain greatly favored  $\text{H}_2\text{O}_2$  partition and diffusion across the membrane. Thus, lipid membranes show different permeability to  $\text{H}_2\text{O}_2$ , depending on the composition, but what happens in the more complex intact RBC membrane?

### Permeability of human RBC membrane to $\text{H}_2\text{O}_2$

The enzyme latency assay was also used to determine the  $P_m$  to  $\text{H}_2\text{O}_2$  in human RBCs. To comply with the requirements of the technique, the experiments were performed with very low cell densities, using the  $\text{HbO}_2$  concentration in the sample as a reference, and with high  $\text{H}_2\text{O}_2$  concentrations (10 mM). Given

these conditions, catalase is the main enzyme involved in  $\text{H}_2\text{O}_2$  decomposition (Fig. S1), as it was demonstrated previously (9). As shown in Figure 3, the rate of  $\text{H}_2\text{O}_2$  decomposition is higher for lysed RBCs than for intact RBCs. The observed rate constant for the disappearance of  $\text{H}_2\text{O}_2$  in lysed RBCs was 4.3 times greater than in intact RBCs, showing that as observed with liposomes, the RBC membrane imposes a barrier to  $\text{H}_2\text{O}_2$  delaying its diffusion and decomposition by the intracellular catalase.

The results obtained with this method led to a  $P_m$  of  $(1.6 \pm 0.3) \times 10^{-3} \text{ cm s}^{-1}$  at 37 °C. Similar experiments performed using 50  $\mu\text{M}$   $\text{H}_2\text{O}_2$ , drawing aliquots in time and quantifying with horseradish peroxidase and *p*-hydroxyphenylacetic acid, yielded very similar  $P_m$  (Fig. S2), validating the experiments performed with 10 mM  $\text{H}_2\text{O}_2$ . To consider the potential contribution of the unstirred layer (USL), which often confounds permeability measurements, especially in larger vesicles such as RBCs (38), we calculated the permeability of an USL 4  $\mu\text{m}$  thick, considering  $D_{\text{H}_2\text{O}_2} = 1.43 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$  (25 °C) (39). The permeability of this layer was calculated to be  $P_{\text{USL}} = 3.6 \times 10^{-2} \text{ cm s}^{-1}$ , which is 36 times greater than  $P_m$  for RBCs at the same temperature ( $(1.0 \pm 0.2) \times 10^{-3} \text{ cm s}^{-1}$ ). Therefore, USL effects were considered to be negligible in these measurements.

The  $P_m$  was also determined at increasing temperatures, up to 40 °C, allowing the estimation of the  $E_a$  of the  $\text{H}_2\text{O}_2$  permeation process (Fig. 3). The calculated value,  $(32 \pm 4) \text{ kJ mol}^{-1}$ , is four times lower than the  $E_a$  associated with simple diffusion through lipid membranes composed of DMPC:DPPG:Chol (4:1:5) and 1.8 times lower than DOPC:POPG:Chol (4:1:5) membranes (Table 2). Although it was expected that the lower  $E_a$  for RBCs would be associated with a higher  $P_m$  (38), this was not observed. Comparison between the liposomes and the RBC membrane can be complicated by the fact that membrane composition is different. In RBC, sphingomyelin that accounts for 25% of the total phospholipids and 14% of the total lipids may alter the behavior of the membrane. Another important difference in the lipids is the asymmetry between inner and outer leaflets (43). Probably the most important factor to explain the difference is the presence of proteins that account for 49% of the total membrane mass and may affect lipid fluidity and packing or offer alternative pathways for  $\text{H}_2\text{O}_2$  diffusion. Altogether, these results are not conclusive about the importance of simple diffusion of  $\text{H}_2\text{O}_2$  across RBC lipid membranes, implying that protein channels may be involved in the transport of  $\text{H}_2\text{O}_2$  across the human RBC membrane.


**Table 2**  
Permeability coefficients and activation energies of permeation for  $\text{H}_2\text{O}_2$  in the different membranes

Liposome or cellular membrane	$P_m$ ( $\text{cm s}^{-1}$ )		$E_a$ ( $\text{kJ mol}^{-1}$ )
	25 °C	37 °C	
DMPC:DPPG:Chol (4:1:5 mole fraction)	$(3.7 \pm 0.5) \times 10^{-5}$	$(4.1 \pm 0.5) \times 10^{-4}$	$130 \pm 20$
DOPC:POPG:Chol (4:1:5 mole fraction)	$(1.3 \pm 0.7) \times 10^{-3}$	$(5.5 \pm 0.3) \times 10^{-3}$	$57 \pm 8$
Intact RBC	$(1.0 \pm 0.2) \times 10^{-3}$	$(1.6 \pm 0.3) \times 10^{-3}$	$32 \pm 4$

$P_m$  is reported as the mean  $\pm$  the standard deviation of at least three independent determinations.

### Role of AQPs in $\text{H}_2\text{O}_2$ membrane diffusion in human RBCs


In other cell types, several AQP isoforms were found to facilitate  $\text{H}_2\text{O}_2$  diffusion through cellular membranes (22). So studies were carried out to assess the role of AQPs in  $\text{H}_2\text{O}_2$  permeability in human RBC. A series of AQP inhibitors were tested, including  $\text{HgCl}_2$  and *p*-chloromercuribenzenesulfonic acid (pCMBS) for AQP1 and phloretin for AQP3, incubating RBCs with each compound to evaluate possible changes in



**Figure 3. Determination of  $P_m$  of RBC membranes to  $\text{H}_2\text{O}_2$ .** A, consumption of 10 mM  $\text{H}_2\text{O}_2$  by intact RBCs at increasing cell densities (0.9–3.5  $\mu\text{M}$   $\text{HbO}_2$ , red tones) and by lysed RBCs (90–350 nM  $\text{HbO}_2$ , blue tones) at 37 °C. B, initial rates of  $\text{H}_2\text{O}_2$  consumption as a function of  $\text{HbO}_2$  concentration for intact (red circles) and lysed (blue squares) cells. The slopes are the respective observed rate constants  $k_{\text{RBC}}$  and  $k_{\text{lys}}$ . The ratio between  $k_{\text{RBC}}$  and  $k_{\text{lys}}$  was used to calculate  $R_{\text{H}_2\text{O}_2}$  and determine  $P_m$  according to Equation 2 ( $n = 3$ , three biological replicates). C, estimation of the  $E_a$  of the  $\text{H}_2\text{O}_2$  permeation process in RBCs,  $E_a = 32 \pm 4 \text{ kJ mol}^{-1}$  ( $n = 3$ ).  $\text{H}_2\text{O}_2$ , hydrogen peroxide;  $\text{HbO}_2$ , oxyhemoglobin; RBC, red blood cell.

$\text{H}_2\text{O}_2$  consumption rates. The canonic (but not specific) AQP1 inhibitor  $\text{HgCl}_2$  resulted in complete inhibition of  $\text{H}_2\text{O}_2$  decomposition by RBCs, but control experiments showed that this was caused by the inhibition of catalase, that is central to

our assay (Fig. 4, A–D), and therefore could not be used. As an alternative, pCMBS was used, and even though it inhibited the transport of water through the RBC membrane as expected, it had no effect on the rate of  $\text{H}_2\text{O}_2$  consumption by intact RBCs



**Figure 4. Effect of aquaporin (AQP) inhibitors on  $\text{H}_2\text{O}_2$  consumption by RBCs.** A and E, water efflux across the RBC membrane (0.5% hematocrit) when mixed with 250 mM sucrose. Changes in cell volume corresponding to control RBCs (black) and RBCs treated with 0.5 mM  $\text{HgCl}_2$  (blue) or 0.5 mM pCMBS (red), in order to inhibit AQP1, were monitored by light scattering measurements. B and F, observed constants ( $k_{\text{water}}$ ) obtained from the exponential fits of time courses resulting from experiments shown, and performed as, in A and E. C, G, and K, consumption of 10 mM  $\text{H}_2\text{O}_2$  by 0.5% hematocrit suspensions of control RBCs (black) and RBCs treated with 0.5 mM  $\text{HgCl}_2$  (blue), 0.5 mM pCMBS (red) or 0.5 mM phloretin (green), an AQP3 inhibitor. D, consumption of 10 mM  $\text{H}_2\text{O}_2$  by solutions of 8 nM catalase untreated (gray) or incubated with 0.3 mM  $\text{HgCl}_2$  (blue). H and L, initial rates of  $\text{H}_2\text{O}_2$  metabolism determined from experiments shown and performed as in G and K. I, glycerol influx across the RBC membrane (0.5% hematocrit) followed by light scattering measurements. Untreated RBCs (black) or RBCs treated with 0.5 mM phloretin (green) were mixed with equal volumes of a 100 mM glycerol solution. J, observed constants ( $k_{\text{glycerol}}$ ) obtained from the nonlinear curve fit of results from experiments performed as in I, corresponding to the decaying phase of the time courses (\*\*\*\* $p < 0.0001$ ).  $\text{H}_2\text{O}_2$ , hydrogen peroxide; pCMBS, *p*-chloromercuribenzenesulfonic acid; RBC, red blood cell.

## Permeability of membranes to hydrogen peroxide

(Fig. 4, E–H). Phloretin inhibited the transport of glycerol, which occurs mostly through AQP3, but had no effect on the rate of  $\text{H}_2\text{O}_2$  decomposition by intact RBCs (Fig. 4, I–L). Considering that the canonical inhibitor of water transport by AQP1,  $\text{HgCl}_2$ , could not be used because of its effect on catalase, and that phloretin may block glycerol transport by AQP3 but not the transport of smaller molecules such as  $\text{H}_2\text{O}_2$ , additional assays were devised to evaluate the importance of these channels in the transport of  $\text{H}_2\text{O}_2$  through the membrane of human RBCs.


To assess the participation of these water channels on the permeability to  $\text{H}_2\text{O}_2$ , human RBCs deficient in AQP1 and AQP3 (Colton-null and GIL-null phenotypes, respectively) were used. These samples have been extensively characterized and found that the Colton-null shows a significantly lower permeability to water, whereas the GIL-null shows normal water permeability but significantly lower permeability to glycerol (44–46). The  $P_m$  to  $\text{H}_2\text{O}_2$  was determined as described previously and compared with wildtype RBCs that had been cryopreserved in the same manner as a control.

As shown in Figure 5, cryopreserved RBC samples retained the permeability barrier to  $\text{H}_2\text{O}_2$ , shown by the higher  $k_{\text{lys}}$  (slope given by *square symbols*) compared with  $k_{\text{RBC}}$  (slope given by *round symbols*). The  $P_m$  for the cryopreserved control RBCs was slightly higher than  $P_m$  determined in fresh RBCs,

but within experimental error. Remarkably, no differences were observed in  $\text{H}_2\text{O}_2$  permeability between control, Colton-null, and GIL-null RBCs. Furthermore, using fresh RBCs, no saturation was observed in  $\text{H}_2\text{O}_2$  consumption rates up to 100 mM  $\text{H}_2\text{O}_2$  (Fig. 5D). Altogether, these results strongly suggest that neither AQP1 nor AQP3 are involved in  $\text{H}_2\text{O}_2$  transport across the human RBC membrane and support that  $\text{H}_2\text{O}_2$  traverses the human RBC membrane by simple diffusion across the lipid fraction or through a still unidentified protein channel.

### Physiological implications of $\text{H}_2\text{O}_2$ diffusion

Because of experimental limitations, most of our results were obtained under nonphysiological conditions, using either very low hematocrit or very high  $\text{H}_2\text{O}_2$  concentration. However, the mathematical model built using known rate constants for the antioxidant enzymes, validated for a wide range of conditions (9) and now refined with the newly obtained  $P_m$ , allows us to explore the otherwise inaccessible physiological conditions of high hematocrit (45%) and low concentration of  $\text{H}_2\text{O}_2$ . The estimated half-life for  $\text{H}_2\text{O}_2$  in these conditions is 35.3 ms (Fig. S3). A higher  $P_m$  would allow for even faster rates of clearance of  $\text{H}_2\text{O}_2$  (Fig. S4). In our experimental conditions with high  $\text{H}_2\text{O}_2$  concentration, Prx2 was rapidly oxidized and




**Figure 5.  $\text{H}_2\text{O}_2$  permeability in control RBCs and RBCs lacking AQP1 (Colton-null) and AQP3 (GIL-null).** A, secondary plot for the  $P_m(\text{H}_2\text{O}_2)$  determination, in control (black) and AQP1-deficient (red) human RBCs. Circles show  $\text{H}_2\text{O}_2$  consumption by intact cells, and squares show consumption by lysed cells. B, secondary plot for the  $P_m(\text{H}_2\text{O}_2)$  determination, in control (black) and AQP3-deficient human RBCs (blue). Circles correspond to intact cells, and squares correspond to lysed cells. C, comparison of  $P_m(\text{H}_2\text{O}_2)$  between control RBCs and RBCs lacking AQP1 (Colton-null) and AQP3 (GIL-null). No statistically significant differences were observed. D, intact (red) and lysed fresh RBCs (blue) were incubated with increasing concentrations of  $\text{H}_2\text{O}_2$ , from 10 to 100 mM, in HBSS solution. The initial rates were determined by stopped flow measuring absorbance at 240 nm for concentrations below 30 mM  $\text{H}_2\text{O}_2$  and 270 nm for concentrations above 50 mM  $\text{H}_2\text{O}_2$  ( $\epsilon(\text{H}_2\text{O}_2) = 7.4 \text{ M}^{-1} \text{ cm}^{-1}$ ). No saturation of  $\text{H}_2\text{O}_2$  transport or consumption by RBCs was observed. AQP1, aquaporin 1; AQP3, aquaporin 3;  $\text{H}_2\text{O}_2$ , hydrogen peroxide; HBSS, Hank's balanced salt solution; RBC, red blood cell.

NADPH depleted (9), and catalase decomposed most  $\text{H}_2\text{O}_2$  (Fig. S1). However, in physiological conditions, Prx2 would remain active, resulting in a much faster decomposition of  $\text{H}_2\text{O}_2$  inside the RBC. As a consequence, a large concentration gradient will be formed across the membrane, estimated to be 1600 times lower in the internal side of the bilayer relative to the external side (Fig. 6). Note that this gradient is much greater than the one formed by catalase alone in the RBC, measured as 4.3 ( $1/R_{\text{H}_2\text{O}_2}$  in Fig. 3). This is because the rate of reaction of Prx2 with  $\text{H}_2\text{O}_2$  is ten times greater and because Prx2 is 20 to 40 times more abundant than catalase (9, 12). The magnitude of the gradient depends inversely on  $P_m$ , so that a 10-fold increase in  $P_m$  would lead to a 10-fold decrease in the gradient (Fig. S4).

## Discussion

The partition of  $\text{H}_2\text{O}_2$  in lipid membranes is difficult to determine because  $\text{H}_2\text{O}_2$  is hydrophilic and thus is excluded from the bilayer. Organic solvents have been used for a long time as substitutes for the estimation of the solubility of molecules in the lipid membrane, in particular octanol and hexadecane (25). Using a double-partition approach, we could determine  $K_D^\circ$  for  $\text{H}_2\text{O}_2$  between water and these solvents and found that  $\text{H}_2\text{O}_2$  is 15 times less soluble in octanol and 122,000 times less soluble in hexadecane than in water at 25 °C. Previous determinations of  $K_D^\circ$  found that  $\text{H}_2\text{O}_2$  was 14 times less soluble in ether than in water (15), whereas attempts using *n*-heptane were reported to be unsuccessful (27). As a comparison, water is 2.4 million times less soluble in hexadecane than in itself and 25 times less soluble in octanol than in itself (32, 47, 48), indicating that  $\text{H}_2\text{O}_2$  is only slightly less hydrophilic than water. Making simplistic assumptions based on solubility



**Figure 6. Concentration gradient of  $\text{H}_2\text{O}_2$  across the membrane.** Extracellular  $\text{H}_2\text{O}_2$  diffuses across the membrane of the RBC and reacts rapidly with Prx2. The fast intracellular decomposition of  $\text{H}_2\text{O}_2$  (given by the very high rate constant  $k = 1 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$  and the high intracellular concentration of Prx2 in the RBC, 400  $\mu\text{M}$ ) and the slightly limited permeability of the membrane to  $\text{H}_2\text{O}_2$  ( $P_m = 1.6 \times 10^{-3} \text{ cm s}^{-1}$ ) allow the formation of a 1600-fold concentration gradient of  $\text{H}_2\text{O}_2$  across the membrane. In contrast, catalase alone generates a 4.3-fold concentration gradient (gray arrow). This large concentration gradient will further promote the net flux of  $\text{H}_2\text{O}_2$  into the RBC, supporting the role of RBCs as efficient sinks of  $\text{H}_2\text{O}_2$  in the vasculature.  $\text{H}_2\text{O}_2$ , hydrogen peroxide; Prx2, peroxiredoxin 2; RBC, red blood cell.

and diffusion, the permeability of membranes composed solely of lipids to  $\text{H}_2\text{O}_2$  was estimated to be  $P_m = 7 \times 10^{-4} \text{ cm s}^{-1}$ .

The Gibbs energy profile constructed with the solubility data suggests that the greatest energy barrier to  $\text{H}_2\text{O}_2$  membrane permeation is located at the middle region of the bilayer (Fig. 1). These results agree with and support several molecular dynamics simulations that provide a more detailed free energy distribution across the bilayer (29). The Gibbs energy was estimated to be  $33 \pm 4 \text{ kJ/mol}$  in a DOPC membrane by molecular dynamics simulation (29), in fair agreement with  $\Delta G^\circ$  between hexadecane and water (29.0 kJ/mol; Table 1).

Although there are many studies on the permeability of cell plasma membranes to  $\text{H}_2\text{O}_2$  (reviewed in Ref. (17)), there are only a few that studied the permeability of membranes composed solely of lipids to  $\text{H}_2\text{O}_2$  (15, 27, 28), and none of them provided a value for  $P_m$ . An interesting observation from those studies was that an increase in membrane fluidity, caused by temperature, changes from the gel to the fluid liquid crystalline phase, or addition of *n*-nonanol resulted in a significant increase in the permeability to  $\text{H}_2\text{O}_2$  (15, 27).

Herein, we could determine  $P_m$  for different membranes and found that it depended greatly on lipid composition (Table 2). Membranes composed of saturated phospholipids and Chol (DMPC:DPPG:Chol 4:1:5) showed a lower  $P_m$  ( $4.1 \times 10^{-4} \text{ cm s}^{-1}$  at 37 °C) than unsaturated phospholipid and Chol membranes (DOPC:POPG:Chol 4:1:5,  $P_m = 5.5 \times 10^{-3} \text{ cm s}^{-1}$  at 37 °C). Because Chol is present at 50%, both membranes are in the liquid ordered state ( $L_O$ ), characterized by a higher order at the acyl chains caused by Chol, while retaining free rotation and two-dimensional fluidity (49–51). However, the condensing effects of Chol are greater on DMPC than on DOPC, evidenced by greater changes in bilayer thickness, apparent area per lipid, and chain order parameters (52). Therefore, membranes containing DOPC:POPG:Chol 4:1:5 will be in a more fluid and less-packed state than the membranes containing the saturated phospholipids, and this would explain the higher  $P_m$ . The higher  $E_a$  of permeation through saturated phospholipid membranes is in agreement with a tighter lipid packing than in unsaturated membranes (Table 2).

In membranes of DOPC:Chol 6:4 at 30 °C,  $P_m$  for water was  $6.8 \times 10^{-3} \text{ cm s}^{-1}$  (53). The permeability of membranes to  $\text{H}_2\text{O}_2$  seems to be slightly lower than the permeability to water, probably because of the smaller size of water, a known factor to make molecules more diffusible through membranes than expected based solely on their  $K_D^\circ$  (25, 54).

The low permeability observed in saturated membranes supports that certain cell membranes may be less permeable to  $\text{H}_2\text{O}_2$  and that channels may be necessary to facilitate the transport of  $\text{H}_2\text{O}_2$ . For instance, yeast membranes lacking functional endogenous AQP homologs were shown to be relatively impermeable to  $\text{H}_2\text{O}_2$ , and this could be reverted by the expression of some AQPs, such as human AQP8 (18). The involvement of AQP8 in mammalian cell transport of  $\text{H}_2\text{O}_2$  was later confirmed (21, 55). AQP3 was also found to facilitate the transport of  $\text{H}_2\text{O}_2$  in mammalian cells (19, 20). In the case

## Permeability of membranes to hydrogen peroxide

of human AQP1, several studies have indicated that it would not facilitate H<sub>2</sub>O<sub>2</sub> transport (18, 19), but this position has been challenged and thus remains in debate (23, 24).

The permeability of the human RBC membranes to H<sub>2</sub>O<sub>2</sub> was found to be  $1.0 \times 10^{-3} \text{ cm s}^{-1}$  and  $1.6 \times 10^{-3} \text{ cm s}^{-1}$ , at 25 and 37 °C, respectively. The measured  $P_m$  in RBC indicates that the membrane will limit the diffusion rate of H<sub>2</sub>O<sub>2</sub> into the cytosol by a factor of  $3 \times 10^4$  (relative to an equally thick layer of water). However, the very fast reaction of H<sub>2</sub>O<sub>2</sub> with Prx2 inside the RBC will allow for an efficient and rapid detoxification of H<sub>2</sub>O<sub>2</sub> (9). A large concentration gradient will be formed across the membrane leading to a net flux of H<sub>2</sub>O<sub>2</sub> from the extracellular space into the RBC (Fig. 6). As a result, under physiological conditions, very little H<sub>2</sub>O<sub>2</sub> that diffuses into the RBC will be able to escape, and RBCs will act as efficient sinks of H<sub>2</sub>O<sub>2</sub> in the vascular space.

In comparison with RBCs from other organisms,  $P_m$  for human RBCs is lower than the one reported for rat RBCs,  $1.2 \times 10^{-2} \text{ cm s}^{-1}$ , at 30 °C (15), but similar to the one reported for horse RBCs,  $6 \times 10^{-4} \text{ cm s}^{-1}$ , at 20 °C (16). The difference in the permeability of different RBCs to H<sub>2</sub>O<sub>2</sub> could be caused by differences in membrane lipid composition and/or different membrane protein contribution. Other cellular and organelle membranes show values between  $4 \times 10^{-4}$  and  $1.6 \times 10^{-3} \text{ cm s}^{-1}$  at 37 °C (17, 56), thus the human RBC is in the high range but is not much more permeable to H<sub>2</sub>O<sub>2</sub> than other cells. The permeability of human RBCs is higher than the permeability of DMPC:DPPG:Chol 4:1:5 membranes but lower than the permeability of DOPC:POPG:Chol 4:1:5 membranes (Table 2). The difference in  $P_m$  could occur because the different composition of lipids modifies the permeability of the membrane, the presence of embedded transmembrane proteins in the RBC membrane that modifies the physical properties of the lipids, or because some transmembrane protein facilitated H<sub>2</sub>O<sub>2</sub> diffusion across the membrane. The  $E_a$  of the permeation process ( $32 \text{ kJ mol}^{-1}$ ) is lower for RBCs than for liposomes (Table 2), and it is in between the values for diffusional and osmotic permeability of regular and AQP1-null human RBCs to water (19 and  $43 \text{ kJ mol}^{-1}$ , respectively (57)), suggesting the involvement of a transporter protein.

The main suspects in facilitating H<sub>2</sub>O<sub>2</sub> transport across the RBC membrane were the AQPs. AQP1 is an abundant protein in the membrane of human RBCs. The number of AQP1 in RBC has been reported in  $6$  to  $14 \times 10^4$  copies (11, 58). Although it is not an essential protein for cell viability, AQP1 is the most important protein in water transport in the RBCs (57). However, our results on H<sub>2</sub>O<sub>2</sub> permeability using AQP1-deficient RBCs (Colton-null) showed no difference to normal human RBCs, suggesting that this channel does not facilitate H<sub>2</sub>O<sub>2</sub> transport in human RBCs.

AQP3 has been repeatedly reported to allow H<sub>2</sub>O<sub>2</sub> transport across the membrane (19, 20), and this property is probably explained by its larger pore that also transports glycerol. Herein, we did not observe differences between regular and

AQP3-deficient human RBCs (GIL-null). The expression level of AQP3 has recently been reported to be 1700 copies per cell, 30 to 100 times less abundant than AQP1 (11). Nonetheless, if H<sub>2</sub>O<sub>2</sub> transport was exclusively supported by AQP3, as described for glycerol, GIL-null RBCs should have been much less permeable than regular RBCs. Considering that no effect on H<sub>2</sub>O<sub>2</sub> was observed, AQP3 is not an important route for its transport across RBC membranes.

It cannot be discarded that other proteins may be involved in facilitating H<sub>2</sub>O<sub>2</sub> transport across the RBC membrane. An interesting candidate is the urea transporter UT-B, which has a unit permeability to water per channel similar to that of AQP1 (44), and is present at  $14$  to  $26 \times 10^3$  copies per cell (11).

Considering that the  $P_m$  for H<sub>2</sub>O<sub>2</sub> across RBC membranes is similar to that of lipid-only liposomes (Table 2); is similar to the membrane osmotic permeability to water in AQP1-null RBCs ( $P_f = 2\text{--}3 \times 10^{-3} \text{ cm s}^{-1}$  at 26 and 20 °C) (57, 59); is similar in regular, AQP1-null, and AQP3-null cells; does not show saturation; strongly suggests that the main mechanism of H<sub>2</sub>O<sub>2</sub> permeation across RBC membranes is simple diffusion across the lipid fraction.

Although in some other cell types, specific AQPs have been shown to facilitate the diffusion of H<sub>2</sub>O<sub>2</sub> (peroxiporins), our results suggest that AQPs are not the only possible routes for H<sub>2</sub>O<sub>2</sub> diffusion into cells. Depending on the cell type (and likely organelle), the lipid fraction of the membrane may be sufficiently permeable to H<sub>2</sub>O<sub>2</sub>, or still unidentified membrane proteins may be involved in facilitating H<sub>2</sub>O<sub>2</sub> transport across cellular membranes.

## Conclusions

In summary, the present study revealed several aspects of H<sub>2</sub>O<sub>2</sub> interaction with membranes: (1) the solubility of H<sub>2</sub>O<sub>2</sub> in the solvents octanol and hexadecane, which resemble the membrane interior, is 15 and 122,000 times lower than in water, confirming that H<sub>2</sub>O<sub>2</sub> will face a large thermodynamic barrier when diffusing across lipid membranes; (2) the permeability of phospholipid:Chol liposome membranes to H<sub>2</sub>O<sub>2</sub> depends on the composition of the lipids and increases with acyl-chain unsaturation; (3) the permeability of RBC at 37 °C is  $1.6 \times 10^{-3} \text{ cm s}^{-1}$ , which is  $3 \times 10^4$  times lower than an equally thick layer of water; (4) the fast reaction of H<sub>2</sub>O<sub>2</sub> with cytosolic Prx2 results in the formation of a 1600-fold concentration gradient of H<sub>2</sub>O<sub>2</sub> across the membrane; (5) the estimated half-life of H<sub>2</sub>O<sub>2</sub> in blood under physiological conditions is 34.5 ms; and (6) AQPs are not involved in facilitating H<sub>2</sub>O<sub>2</sub> diffusion across the membrane of RBCs, and diffusion occurs likely through the lipid fraction or through a still unidentified membrane protein.

## Experimental procedures

### Materials

Chemical reagents were obtained from Sigma, Applichem, and Acros Organics. Lipids were obtained from Avanti Polar



Lipids and Larodan. Work solutions of H<sub>2</sub>O<sub>2</sub> were prepared daily and quantified by spectrophotometry,  $\epsilon(240 \text{ nm}) = 39.4 \text{ M}^{-1} \text{ cm}^{-1}$  (60).

The studies using human blood were conducted in accordance with the Declaration of Helsinki. Blood was obtained from volunteer donors after informed consent at the Cátedra y Departamento de Medicina Transfusional del Hospital de Clínicas, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay. The research protocol was approved by the Hospital's Ethics Committee. Packed RBCs were obtained by standard techniques without leukoreduction, as described before (61). For AQP1-deficient and AQP3-deficient RBC experiments, control, Colton-null, and GIL-null RBCs were obtained from cryopreserved samples from the Rare Blood Collection at the Centre National de Référence pour les Groupes Sanguins de l'Institut National de la Transfusion Sanguine, Paris, France. Spectrophotometric measurements were performed in a Varian Cary 50 (Agilent) spectrophotometer in wildtype experiments. In AQP1-deficient and AQP3-deficient RBC experiments, a NanoDrop 2000c spectrophotometer (Thermo Fisher Scientific) was used for simple spectrophotometric reads, and an SFM400 Stopped Flow (Bio-Logic) was used for continuous absorbance measurements.

### H<sub>2</sub>O<sub>2</sub> partition experiments

The solubility of H<sub>2</sub>O<sub>2</sub> in organic solvents is very low because H<sub>2</sub>O<sub>2</sub> is hydrophilic. Furthermore, most methods for H<sub>2</sub>O<sub>2</sub> quantification are based on water. Therefore, we devised a double-partition method involving a first step of equilibration of H<sub>2</sub>O<sub>2</sub> between water (w1) and organic solvent (os1), careful extraction of the organic phase to prevent carrying any water, and then a second step where this organic phase (os2) was equilibrated with new water (w2). Every volume was accurately determined (V<sub>os1</sub>, V<sub>w1</sub>, V<sub>os2</sub>, and V<sub>w2</sub>). Note that V<sub>os2</sub> was approximately half that of V<sub>os1</sub> to avoid any contaminating water. The water used was saturated with the organic solvent of interest, and the organic solvents were saturated with water to prevent artificial increases in the solubility of H<sub>2</sub>O<sub>2</sub>. The equilibration of a known amount of H<sub>2</sub>O<sub>2</sub> proceeded at controlled temperature in a water bath with frequent mixing for 30 min. The phases were then separated by centrifugation, and a fraction of the organic solvent (on top) was removed and placed in a new container with water, and the equilibration was repeated. The concentration of H<sub>2</sub>O<sub>2</sub> was quantified both in the initial and final water phases.

The following rationale was used to calculate the  $K_D^\circ$  of H<sub>2</sub>O<sub>2</sub> between the organic phase and water. First,  $K_D^\circ$  is the same in both steps, while the concentrations in each phase change.

$$K_D^\circ = [\text{H}_2\text{O}_2]_{\text{os1}} / [\text{H}_2\text{O}_2]_{\text{w1}} \quad (3)$$

$$K_D^\circ = [\text{H}_2\text{O}_2]_{\text{os2}} / [\text{H}_2\text{O}_2]_{\text{w2}} \quad (4)$$

Because of mass conservation, the total number of moles in the second equilibrium (n<sub>T2</sub>) is:

$$n_{T2} = [\text{H}_2\text{O}_2]_{\text{w2}} V_{\text{w2}} + [\text{H}_2\text{O}_2]_{\text{os2}} V_{\text{os2}} \quad (5)$$

that can be expressed as

$$n_{T2} = [\text{H}_2\text{O}_2]_{\text{w2}} (V_{\text{w2}} + K_D^\circ V_{\text{os2}}) \quad (6)$$

With this information, we can calculate the concentration in the organic solvent in the first equilibrium:

$$[\text{H}_2\text{O}_2]_{\text{os1}} = n_{T2} / V_{\text{os2}} \quad (7)$$

So that we can replace in Equation 3 and rearrange to obtain:

$$K_D^\circ = [\text{H}_2\text{O}_2]_{\text{w2}} V_{\text{w2}} / (([\text{H}_2\text{O}_2]_{\text{w1}} - [\text{H}_2\text{O}_2]_{\text{w2}}) V_{\text{os2}}) \quad (8)$$

The quantification of H<sub>2</sub>O<sub>2</sub> was done using horseradish peroxidase (0.05 U/ml) and 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (0.1 mM) in 0.1 M potassium phosphate buffer at pH 5, measuring the 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) radical cation at 734 nm by spectrophotometry (62). Calibration curves were constructed using H<sub>2</sub>O<sub>2</sub> quantified at 240 nm (see aforementioned). The response was linear up to 40 μM. Different conditions were assayed until optimal initial concentrations in w1 for octanol and hexadecane were found to be 5 mM and 2 M H<sub>2</sub>O<sub>2</sub>, respectively.

### Thermodynamics of partition

The Gibbs energy of partition ( $\Delta G^\circ$ ), indicating the energy required for the transfer of H<sub>2</sub>O<sub>2</sub> from water to the organic solvent, was calculated from  $K_D^\circ$ :

$$\Delta G^\circ = -RT \ln(K_D^\circ) \quad (9)$$

where  $R$  is the universal gas constant (8.314 J mol<sup>-1</sup> K<sup>-1</sup>) and  $T$  is the absolute temperature.

The enthalpy and entropy of the partition ( $\Delta H^\circ$  and  $\Delta S^\circ$ ) were calculated from  $K_D^\circ$  at 25 and 37 °C, using the van't Hoff equation:

$$\ln(K_D^\circ) = -\frac{H^\circ}{R} \frac{1}{T} + \frac{S^\circ}{R} \quad (10)$$

### Preparation of catalase-encapsulated liposomes

The desired final composition of the membrane was made by mixing chloroform solutions of the lipids and then drying by nitrogen stream and then vacuum for 2 h. Typical preparations contained 10 mg of lipids and were composed of either DMPC:DPPG:Chol (4:1:5 mole fraction) or DOPC:POPG:Chol (4:1:5 mole fraction).

## Permeability of membranes to hydrogen peroxide

The dried lipids were then incubated with a work solution containing 2 mg/ml catalase in 50 mM sodium phosphate at pH 7 buffer. Catalase solution was previously filtered through 0.22  $\mu\text{m}$  pore filters (polyvinylidene fluoride; Millipore). The lipids were allowed to hydrate in this solution for 30 min with frequent mixing. Liposomes were then prepared by extrusion through 200 nm or 1000 nm nucleopore membranes (Whatman) 15 times using a syringe extruder (Avanti). To separate the catalase-encapsulated liposomes from free catalase, a Superdex 200 (10/300 GL; GE Healthcare) column coupled to an HPLC (Agilent; 1260), using 50 mM sodium phosphate at pH 7 at 0.5 ml/min as the mobile phase, was used. Because of their large size, liposomes elute very early (17 min), whereas catalase is retarded and elutes later (30 min). Liposomes were collected and used for permeability experiments. The size of the liposomes was consistent with the nominal pore value of the nucleopore filter membranes to within 5%, as confirmed by dynamic light scattering (Brookhaven Instruments).

### Determination of permeability coefficients of lipid membranes

The calculation of the  $P_m$  is based on the concept of enzyme latency (41). In some cases, enzymes encapsulated by membranes show a lower activity than enzymes free in solution, particularly if the membrane is partially permeable to the substrate. The diffusion of the substrate is slowed, and a concentration gradient forms across the membrane that is observed as a different enzyme activity. In steady state, the ratio of membrane-encapsulated and free enzyme activity ( $R_{\text{H}_2\text{O}_2}$ ) will yield the concentration gradient. For  $\text{H}_2\text{O}_2$ , the catalase activity is measured by following the rate of  $\text{H}_2\text{O}_2$  (10 mM) decomposition by spectrophotometry at 240 nm, by liposome-encapsulated and disrupted liposome catalase (16, 41). Liposomes containing catalase were disrupted by filtration through 30 nm nucleopore filters (Whatman) 11 times in a syringe mini extruder (Avanti). Complete equilibration of catalase between external and internal volumes was confirmed by measuring catalase activity at different cycles of extrusion, which showed that seven cycles already yielded maximal catalase activity. Initial rates of  $\text{H}_2\text{O}_2$  decomposition were obtained from linear regression and plotted as a function of liposome volume. The slopes of these secondary plots (named  $k_{\text{lipo}}$  and  $k_{\text{dis}}$  for intact and disrupted liposomes, respectively) were used to calculate  $R_{\text{H}_2\text{O}_2}$  ( $k_{\text{lipo}}/k_{\text{dis}}$ ) and then  $P_m$  according to Equation 2.  $k_{\text{catalase}}$  was determined by measuring the rate of decomposition of  $\text{H}_2\text{O}_2$  by different dilutions of the catalase work solution and then extrapolated to the concentration in the work solution. This  $k_{\text{catalase}}$  was determined individually for each liposome preparation.

### Estimation of activation energy for $\text{H}_2\text{O}_2$ permeation

$P_m$  was determined at different temperatures between 10 and 40  $^\circ\text{C}$  following the previously described protocol. The obtained values were used to construct a  $\ln(P_m)$  versus  $T^{-1}$

plot, from which the  $E_a$  was calculated as shown in Equation 11:

$$\ln(P_m) = \ln(A) - \frac{E_a}{R} \left( \frac{1}{T} \right) \quad (11)$$

where  $A$  is the pre-exponential factor and  $R$  is the universal gas constant. The  $E_a$  for catalase is very low (2.5 kJ/mol, (63)), and the  $E_a$  for  $P_m$  was not corrected for its contribution.

### RBC preparation

Before each experiment, the RBCs were washed three times in Hank's balanced salt solution (137 mM NaCl, 5.4 mM KCl, 0.25 mM  $\text{Na}_2\text{HPO}_4$ , 1 g  $\text{l}^{-1}$  glucose, 0.44 mM  $\text{KH}_2\text{PO}_4$ , 1.3 mM  $\text{CaCl}_2$ , 1.0 mM  $\text{MgSO}_4$  and 4.2 mM  $\text{NaHCO}_3$ , and pH 7.4) by centrifugations at 900g for 4 min at room temperature. After that, the cells were resuspended in the same solution and diluted until the hematocrit was approximately 0.06%. The experiments with RBCs were performed using freshly obtained blood, on the same day of the experiment. The experiments with RBCs lacking AQP1 or AQP3 were performed after recovering the frozen samples and compared with cryopreserved normal RBCs.

### Determination of $\text{H}_2\text{O}_2$ permeability coefficient of RBCs

Suspensions of intact RBCs with increasing hematocrits, obtained *via* dilution of the stock, were mixed with 10 mM  $\text{H}_2\text{O}_2$  in Hank's balanced salt solution.  $\text{H}_2\text{O}_2$  consumption was then measured spectrophotometrically at 240 nm for 1 min at 37  $^\circ\text{C}$ . The same procedure was performed using lysed cells, generated by freezing part of the original stock of RBCs. Initial rates were obtained from linear regression and plotted as a function of  $\text{HbO}_2$  concentration, determined through absorbance measurements at 577 nm ( $\epsilon(\text{HbO}_2) = 15 \text{ mM}^{-1} \text{ cm}^{-1}$ , (64)). The slopes of these secondary plots (named  $k_{\text{RBC}}$  and  $k_{\text{lys}}$  for intact and lysed RBCs, respectively) were used to calculate  $R_{\text{H}_2\text{O}_2}$  and then  $P_m$  following Equation 10 (41). Hemolysis was very low in these experiments. Free hemoglobin after treatment with  $\text{H}_2\text{O}_2$  was the same as the control with buffer and less than 2% of the equivalent lysed RBCs. Considering that intact RBCs had 23% of the catalase activity compared with lysed RBCs, more than ten times than that expected from the contribution of hemolysis, it can be ascertained that in intact RBC experiments, most of the catalase activities were derived from RBC-encapsulated catalase. Because some of the hemolysis may also occur during the centrifugation to sediment RBCs, no corrections for hemolysis were introduced.

In the case of RBCs,  $k_{\text{catalase}}$  is the pseudo-first-order constant for  $\text{H}_2\text{O}_2$  removal by catalase inside the RBC (determined by extrapolation of the  $k_{\text{lys}}$  value for a concentration of 20 mM  $\text{HbO}_2$ ),  $R_{\text{H}_2\text{O}_2}$  represents the  $k_{\text{RBC}}/k_{\text{lys}}$  ratio,  $A$  is the surface area ( $1.4 \times 10^{-6} \text{ cm}^2$ ) and  $V$  the volume ( $9 \times 10^{-11} \text{ cm}^3$ ) of the RBC (65, 66). Determinations of  $P_m$  were done using three biological replicates, each measured three

separate times ( $n = 9$ ) in order to determine  $P_m$  in basal conditions (37 °C, pH 7.4). Mathematical simulations showed that under these experimental conditions, catalase was effectively the most important enzymatic system decomposing  $H_2O_2$  (>97%; Fig. S1).

### Water permeability and inhibition of AQP1

RBC suspensions at 0.5% hematocrit were incubated with 0.5 mM  $HgCl_2$  or pCMBS in PBS (137 mM NaCl, 2.7 mM KCl, 8 mM  $Na_2HPO_4$ , and 2 mM  $KH_2PO_4$ ) for 30 min at room temperature. Inhibition of water passage through AQP1 by these compounds was checked using an SX Stopped-Flow Spectrometer (Applied Photophysics) to follow changes in RBC volume after mixing them with an equal volume of a hypertonic solution of 250 mM sucrose at 10 °C (67). Water efflux was recorded by 90° light scattering measurements of 2 s of duration, and the resulting time courses were fitted to an exponential model. The observed constants ( $k_{water}$ ) were then contrasted with those obtained in assays involving untreated RBCs. The effect of  $HgCl_2$  and pCMBS in  $H_2O_2$  metabolization was evaluated by taking a 40  $\mu$ l aliquot of the 0.5% hematocrit stock, treated or untreated with each inhibitor, and adding 10 mM  $H_2O_2$  in a final volume of 1 ml.  $H_2O_2$  removal was monitored by absorbance measurements at 240 nm, allowing the calculation of initial rates of decomposition.

### Glycerol permeability and inhibition of AQP3

Phloretin was tested and used as an AQP3 inhibitor. RBC suspensions of 0.5% hematocrit were incubated in hypotonic 0.7 $\times$  PBS for 30 min at room temperature. Afterward, a fraction of the sample was treated with 0.5 mM phloretin for another 30 min, whereas the rest remained untreated. Control or treated RBCs were then mixed in equal volumes with a solution of 100 mM glycerol in hypotonic buffer using an SX Stopped-Flow Spectrometer (Applied Photophysics) (68). Changes in cell volume corresponding to the efflux of water and subsequent influx of glycerol were followed by light scattering measurements for 40 s. The resulting time courses were fitted to a double exponential function with two observed constants corresponding to the ascending and descending phases, the latter ( $k_{glycerol}$ ) being used to evaluate the effect of phloretin in glycerol transport (68). To evaluate the effect in  $H_2O_2$  consumption, the cells (0.5% hematocrit) were incubated with 0.5 mM phloretin in 1 $\times$  PBS. The reaction with  $H_2O_2$  was studied in the same conditions as described previously.

### Statistical analysis

Data were analyzed using GraphPad Prism 6 (GraphPad Software, Inc). Statistical analyses were performed by one-way ANOVA and Dunnett's post hoc test to perform multiple comparison tests. Differences with  $p < 0.05$  were considered statistically significant.

### Data availability

All data are contained within the article. Copasi files are to be shared upon request to Matías Möller, [mmoller@fcien.edu.uy](mailto:mmoller@fcien.edu.uy).

*Supporting information*—This article contains supporting information (9, 16, 17, 41, 42, 61, 69–72).

*Acknowledgments*—We acknowledge Dr Thierry Peyrard, head of the Centre National de Référence pour les Groupes Sanguins, at the Institut National de la Transfusion Sanguine, for authorizing the thawing of rare blood samples GIL-null and Colton-null and Dr Eduardo Mendez, Instituto de Química Biológica, Universidad de la República, for useful discussions.

*Author contributions*—L. T. and M. N. M. conceptualization; L. T. and M. N. M. methodology; I. M.-C. and M. A. O. formal analysis; F. O., A. C. L., D. S., C. A., L. T., and M. N. M. investigation; I. R.-G., I. M.-C., and M. A. O. resources; F. O., A. C. L., L. T., and M. N. M. writing—original draft; F. O., A. C. L., I. R.-G., I. M.-C., M. A. O., A. D., L. T., and M. N. M. writing—review & editing; L. T. and M. N. M. supervision; A. D., L. T., and M. N. M. project administration; A. D., L. T., and M. N. M. funding acquisition.

*Funding and additional information*—The authors acknowledge financial support from *Agencia Nacional de Investigación e Innovación*, Uruguay (grants FCE\_2017\_136043 to L. T. and M. N. M. and FMV\_2019\_155597 to L. T. and I. R.-G., and a fellowship to A. C. L.) and to *Comisión Sectorial de Investigación Científica* (grants *Iniciación a la Investigación Científica* to F. O. and C. A.; and *Programa Grupos I+D*\_2014\_C632-348 and \_2018\_47 to A. D.). F. O. received a scholarship from *Comisión Académica de Posgrados*, *Universidad de la República*.

*Conflict of interest*—The authors declare that they have no conflicts of interest with the contents of this article.

*Abbreviations*—The abbreviations used are: AQP1, aquaporin 1; AQP3, aquaporin 3; AQP8, aquaporin 8; Chol, cholesterol; DMPC, dimiristoylphosphatidylcholine; DOPC, dioleoylphosphatidylcholine; DPPG, dipalmitoylphosphatidylglycerol; HbO<sub>2</sub>, oxyhemoglobin;  $H_2O_2$ , hydrogen peroxide; pCMBS, *p*-chloromercuribenzenesulfonic acid; POPG, phosphatidylglycerol; Prx2, peroxiredoxin 2; RBC, red blood cell; USL, unstirred layer.

### References

- Bretón-Romero, R., and Lamas, S. (2014) Hydrogen peroxide signaling in vascular endothelial cells. *Redox Biol.* **2**, 529–534
- Sobotta, M. C., Liou, W., Stöcker, S., Talwar, D., Oehler, M., Ruppert, T., Scharf, A. N., and Dick, T. P. (2015) Peroxiredoxin-2 and STAT3 form a redox relay for  $H_2O_2$  signaling. *Nat. Chem. Biol.* **11**, 64–70
- Stöcker, S., Van Laer, K., Mijuskovic, A., and Dick, T. P. (2018) The conundrum of hydrogen peroxide signaling and the emerging role of peroxiredoxins as redox relay hubs. *Antioxid. Redox Signal.* **28**, 558–573
- Sato, A., Sakuma, I., and Gutterman, D. D. (2003) Mechanism of dilation to reactive oxygen species in human coronary arterioles. *Am. J. Physiol. Heart Circ. Physiol.* **285**, H2345–H2354
- Thengchaisri, N., and Kuo, L. (2003) Hydrogen peroxide induces endothelium-dependent and-independent coronary arteriolar dilation: Role of cyclooxygenase and potassium channels. *Am. J. Physiol. Heart Circ. Physiol.* **285**, H2255–H2263

## Permeability of membranes to hydrogen peroxide

- Abid, M. R., Kachra, Z., Spokes, K. C., and Aird, W. C. (2000) NADPH oxidase activity is required for endothelial cell proliferation and migration. *FEBS Lett.* **486**, 252–256
- Stone, J. R., and Collins, T. (2002) The role of hydrogen peroxide in endothelial proliferative responses. *Endothelium* **9**, 231–238
- Forman, H. J., Bernardo, A., and Davies, K. J. (2016) What is the concentration of hydrogen peroxide in blood and plasma? *Arch. Biochem. Biophys.* **603**, 48–53
- Orrico, F., Möller, M. N., Cassina, A., Denicola, A., and Thomson, L. (2018) Kinetic and stoichiometric constraints determine the pathway of H<sub>2</sub>O<sub>2</sub> consumption by red blood cells. *Free Radic. Biol. Med.* **121**, 231–239
- Moore, R. B., Mankad, M. V., Shriver, S. K., Mankad, V. N., and Plishker, G. A. (1991) Reconstitution of Ca (2+)-dependent K<sup>+</sup> transport in erythrocyte membrane vesicles requires a cytoplasmic protein. *J. Biol. Chem.* **266**, 18964–18968
- Bryk, A. H., and Wiśniewski, J. R. (2017) Quantitative analysis of human red blood cell proteome. *J. Proteome Res.* **16**, 2752–2761
- Manta, B., Hugo, M., Ortiz, C., Ferrer-Sueta, G., Trujillo, M., and Denicola, A. (2009) The peroxidase and peroxynitrite reductase activity of human erythrocyte peroxiredoxin 2. *Arch. Biochem. Biophys.* **484**, 146–154
- Low, F. M., Hampton, M. B., Peskin, A. V., and Winterbourn, C. C. (2006) Peroxiredoxin 2 functions as a noncatalytic scavenger of low-level hydrogen peroxide in the erythrocyte. *Blood* **109**, 2611–2617
- Benfeitás, R., Selvaggio, G., Antunes, F., Coelho, P. M., and Salvador, A. (2014) Hydrogen peroxide metabolism and sensing in human erythrocytes: A validated kinetic model and reappraisal of the role of peroxiredoxin II. *Free Radic. Biol. Med.* **74**, 35–49
- Mathai, J. C., and Sitaramam, V. (1994) Stretch sensitivity of transmembrane mobility of hydrogen peroxide through voids in the bilayer. Role of cardiolipin. *J. Biol. Chem.* **269**, 17784–17793
- Nicholls, P. (1965) Activity of catalase in the red cell. *Biochim. Biophys. Acta* **99**, 286–297
- Möller, M. N., Cuevasanta, E., Orrico, F., Lopez, A. C., Thomson, L., and Denicola, A. (2019) Diffusion and transport of reactive species across cell membranes. In *Bioactive Lipids in Health and Disease*, Springer, Cham: 3–19
- Bienert, G. P., Möller, A. L., Kristiansen, K. A., Schulz, A., Møller, I. M., Schjoerring, J. K., and Jahn, T. P. (2007) Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J. Biol. Chem.* **282**, 1183–1192
- Miller, E. W., Dickinson, B. C., and Chang, C. J. (2010) Aquaporin-3 mediates hydrogen peroxide uptake to regulate downstream intracellular signaling. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 15681–15686
- Hara-Chikuma, M., Chikuma, S., Sugiyama, Y., Kabashima, K., Verkman, A. S., Inoue, S., and Miyachi, Y. (2012) Chemokine-dependent T cell migration requires aquaporin-3-mediated hydrogen peroxide uptake. *J. Exp. Med.* **209**, 1743–1752
- Bertolotti, M., Bestetti, S., García-Manteiga, J. M., Medraño-Fernandez, I., Dal Mas, A., Malosio, M. L., and Sitia, R. (2013) Tyrosine kinase signal modulation: A matter of H<sub>2</sub>O<sub>2</sub> membrane permeability? *Antioxid. Redox Signal.* **19**, 1447–1451
- Bienert, G. P., and Chaumont, F. (2014) Aquaporin-facilitated transmembrane diffusion of hydrogen peroxide. *Biochim. Biophys. Acta* **1840**, 1596–1604
- Almasalmeh, A., Krenc, D., Wu, B., and Beitz, E. (2014) Structural determinants of the hydrogen peroxide permeability of aquaporins. *FEBS J.* **281**, 647–656
- Montiel, V., Bella, R., Michel, L. Y., Esfahani, H., De Mulder, D., Robinson, E. L., Deglasse, J.-P., Tiburcy, M., Chow, P. H., and Jonas, J.-C. (2020) Inhibition of aquaporin-1 prevents myocardial remodeling by blocking the transmembrane transport of hydrogen peroxide. *Sci. Transl. Med.* **12**, eaay2176
- Walter, A., and Gutknecht, J. (1986) Permeability of small nonelectrolytes through lipid bilayer membranes. *J. Membr. Biol.* **90**, 207–217
- Missner, A., and Pohl, P. (2009) 110 Years of the Meyer–Overton rule: Predicting membrane permeability of gases and other small compounds. *ChemPhysChem* **10**, 1405–1414
- Abuin, E., Lissi, E., and Ahumada, M. (2012) Diffusion of hydrogen peroxide across DPPC large unilamellar liposomes. *Chem. Phys. Lipids* **165**, 656–661
- Wang, H., Schoebel, S., Schmitz, F., Dong, H., and Hedfalk, K. (2020) Characterization of aquaporin-driven hydrogen peroxide transport. *Biochim. Biophys. Acta Biomembr.* **1862**, 183065
- Cordeiro, R. M. (2014) Reactive oxygen species at phospholipid bilayers: Distribution, mobility and permeation. *Biochim. Biophys. Acta Biomembr.* **1838**, 438–444
- Yusupov, M., Yan, D., Cordeiro, R. M., and Bogaerts, A. (2018) Atomic scale simulation of H<sub>2</sub>O<sub>2</sub> permeation through aquaporin: Toward the understanding of plasma cancer treatment. *J. Phys. D Appl. Phys.* **51**, 125401
- Razzokov, J., Yusupov, M., Cordeiro, R. M., and Bogaerts, A. (2018) Atomic scale understanding of the permeation of plasma species across native and oxidized membranes. *J. Phys. D Appl. Phys.* **51**, 365203
- Finkelstein, A. (1976) Water and nonelectrolyte permeability of lipid bilayer membranes. *J. Gen. Physiol.* **68**, 127–135
- Xiang, T.-X., and Anderson, B. D. (1998) Influence of chain ordering on the selectivity of dipalmitoylphosphatidylcholine bilayer membranes for permeant size and shape. *Biophys. J.* **75**, 2658–2671
- Marsh, D. (2002) Membrane water-penetration profiles from spin labels. *Eur. Biophys. J.* **31**, 559–562
- Nymeyer, H., and Zhou, H.-X. (2008) A method to determine dielectric constants in nonhomogeneous systems: Application to biological membranes. *Biophys. J.* **94**, 1185–1193
- Stern, H. A., and Feller, S. E. (2003) Calculation of the dielectric permittivity profile for a nonuniform system: Application to a lipid bilayer simulation. *J. Chem. Phys.* **118**, 3401–3412
- Diamond, J. M., and Katz, Y. (1974) Interpretation of nonelectrolyte partition coefficients between dimyristoyl lecithin and water. *J. Membr. Biol.* **17**, 121–154
- Hanneschlaeger, C., Horner, A., and Pohl, P. (2019) Intrinsic membrane permeability to small molecules. *Chem. Rev.* **119**, 5922–5953
- van Stroe-Biezen, S., Everaerts, F., Janssen, L., and Tackens, R. (1993) Diffusion coefficients of oxygen, hydrogen peroxide and glucose in a hydrogel. *Anal. Chim. Acta* **273**, 553–560
- Torchilin, V., and Weissig, V. (2003) *Liposomes: A Practical Approach*, 2nd ed., Oxford University Press, Oxford
- Antunes, F., and Cadenas, E. (2000) Estimation of H<sub>2</sub>O<sub>2</sub> gradients across biomembranes. *FEBS Lett.* **475**, 121–126
- Marinho, H. S., Cyrne, L., Cadenas, E., and Antunes, F. (2013) The cellular steady-state of H<sub>2</sub>O<sub>2</sub>: Latency concepts and gradients. *Methods Enzymol.* **527**, 3–19
- Lorent, J., Levental, K., Ganesan, L., Rivera-Longworth, G., Sezgin, E., Doktorova, M., Lyman, E., and Levental, I. (2020) Plasma membranes are asymmetric in lipid unsaturation, packing and protein shape. *Nat. Chem. Biol.* **16**, 644–652
- Azouzi, S., Gueroult, M., Ripoche, P., Genetet, S., Aronovicz, Y. C., Le Van Kim, C., Etchebest, C., and Mouro-Chanteloup, I. (2013) Energetic and molecular water permeation mechanisms of the human red blood cell urea transporter B. *PLoS One* **8**, e82338
- Preston, G. M., Smith, B. L., Zeidel, M. L., Moulds, J. J., and Agre, P. (1994) Mutations in aquaporin-1 in phenotypically normal humans without functional CHIP water channels. *Science* **265**, 1585–1587
- Roudier, N., Ripoche, P., Gane, P., Le Pennec, P. Y., Daniels, G., Cartron, J.-P., and Bailly, P. (2002) AQP3 deficiency in humans and the molecular basis of a novel blood group system, GIL. *J. Biol. Chem.* **277**, 45854–45859
- Schatzberg, P. (1963) Solubilities of water in several normal alkanes from C<sub>7</sub> to C<sub>16</sub>. *J. Phys. Chem.* **67**, 776–779
- Wolosin, J., and Ginsburg, H. (1975) The permeation of organic acids through lecithin bilayers. Resemblance to diffusion in polymers. *Biochim. Biophys. Acta* **389**, 20–33
- Almeida, P. F., Vaz, W. L., and Thompson, T. (1992) Lateral diffusion in the liquid phases of dimyristoylphosphatidylcholine/cholesterol lipid bilayers: A free volume analysis. *Biochemistry* **31**, 6739–6747
- Nyholm, T. K., Lindroos, D., Westerlund, B., and Slotte, J. P. (2011) Construction of a DOPC/PSM/cholesterol phase diagram based on the fluorescence properties of trans-parinaric acid. *Langmuir* **27**, 8339–8350

51. Mateo, C. R., Acuna, A. U., and Brochon, J.-C. (1995) Liquid-crystalline phases of cholesterol/lipid bilayers as revealed by the fluorescence of trans-parinaric acid. *Biophys. J.* **68**, 978–987
52. Pan, J., Tristram-Nagle, S., and Nagle, J. F. (2009) Effect of cholesterol on structural and mechanical properties of membranes depends on lipid chain saturation. *Phys. Rev. E* **80**, 021931
53. Mathai, J. C., Tristram-Nagle, S., Nagle, J. F., and Zeidel, M. L. (2008) Structural determinants of water permeability through the lipid membrane. *J. Gen. Physiol.* **131**, 69–76
54. Lieb, W. R., and Stein, W. D. (1986) Non-Stokesian nature of transverse diffusion within human red cell membranes. *J. Membr. Biol.* **92**, 111–119
55. Marchissio, M. J., Francés, D. E. A., Carnovale, C. E., and Marinelli, R. A. (2012) Mitochondrial aquaporin-8 knockdown in human hepatoma HepG2 cells causes ROS-induced mitochondrial depolarization and loss of viability. *Toxicol. Appl. Pharmacol.* **264**, 246–254
56. Makino, N., Sasaki, K., Hashida, K., and Sakakura, Y. (2004) A metabolic model describing the H<sub>2</sub>O<sub>2</sub> elimination by mammalian cells including H<sub>2</sub>O<sub>2</sub> permeation through cytoplasmic and peroxisomal membranes: Comparison with experimental data. *Biochim. Biophys. Acta* **1673**, 149–159
57. Mathai, J. C., Mori, S., Smith, B. L., Preston, G. M., Mohandas, N., Collins, M., Van Zijl, P. C., Zeidel, M. L., and Agre, P. (1996) Functional analysis of aquaporin-1 deficient red cells. The Colton-null phenotype. *J. Biol. Chem.* **271**, 1309–1313
58. Burton, N. M., and Bruce, L. J. (2011) Modelling the structure of the red cell membrane. *Biochem. Cell Biol.* **89**, 200–215
59. Roudier, N., Verbavatz, J.-M., Maurel, C., Ripoche, P., and Tacnet, F. (1998) Evidence for the presence of aquaporin-3 in human red blood cells. *J. Biol. Chem.* **273**, 8407–8412
60. Nelson, D. P., and Kiesow, L. A. (1972) Enthalpy of decomposition of hydrogen peroxide by catalase at 25 C (with molar extinction coefficients of H<sub>2</sub>O<sub>2</sub> solutions in the UV). *Anal. Biochem.* **49**, 474–478
61. Amen, F., Machin, A., Touriño, C., Rodríguez, I., Denicola, A., and Thomson, L. (2017) N-acetylcysteine improves the quality of red blood cells stored for transfusion. *Arch. Biochem. Biophys.* **621**, 31–37
62. Cai, H., Liu, X., Zou, J., Xiao, J., Yuan, B., Li, F., and Cheng, Q. (2018) Multi-wavelength spectrophotometric determination of hydrogen peroxide in water with peroxidase-catalyzed oxidation of ABTS. *Chemosphere* **193**, 833–839
63. Beers, R. F., and Sizer, I. W. (1952) A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J. Biol. Chem.* **195**, 133–140
64. Winterbourn, C. C. (1990) [26] Oxidative reactions of hemoglobin. In: Packer, L., Glazer, A. N., eds., *Methods in Enzymology* **186**. Elsevier, San Diego: 265–272
65. Evans, E., and Fung, Y.-C. (1972) Improved measurements of the erythrocyte geometry. *Microvasc. Res.* **4**, 335–347
66. Park, H., Lee, S., Ji, M., Kim, K., Son, Y., Jang, S., and Park, Y. (2016) Measuring cell surface area and deformability of individual human red blood cells over blood storage using quantitative phase imaging. *Sci. Rep.* **6**, 34257
67. Yang, B., Kim, J. K., and Verkman, A. (2006) Comparative efficacy of HgCl<sub>2</sub> with candidate aquaporin-1 inhibitors DMSO, gold, TEA<sup>+</sup> and acetazolamide. *FEBS Lett.* **580**, 6679–6684
68. Rodriguez, R. A., Liang, H., Chen, L. Y., Plascencia-Villa, G., and Perry, G. (2019) Single-channel permeability and glycerol affinity of human aquaporin AQP3. *Biochim. Biophys. Acta Biomembr.* **1861**, 768–775
69. Hoops, S., Sahle, S., Gauges, R., Lee, C., Pahle, J., Simus, N., Singhal, M., Xu, L., Mendes, P., and Kummer, U. (2006) COPASI—A complex pathway simulator. *Bioinformatics* **22**, 3067–3074
70. Winterbourn, C. C. (2013) The biological chemistry of hydrogen peroxide. *Methods Enzymol.* **528**, 3–25
71. Trujillo, M., Alvarez, B., and Radi, R. (2016) One- and two-electron oxidation of thiols: Mechanisms, kinetics and biological fates. *Free Radic. Res.* **50**, 150–171
72. Takebe, G., Yarimizu, J., Saito, Y., Hayashi, T., Nakamura, H., Yodoi, J., Nagasawa, S., and Takahashi, K. (2002) A comparative study on the hydroperoxide and thiol specificity of the glutathione peroxidase family and selenoprotein P. *J. Biol. Chem.* **277**, 41254–41258