

3D models related to the publication: 3D Finite Element Analysis and Geometric Morphometrics of Sloths (Xenarthra, Folivora) Mandibles Show Insights on the Dietary Specializations of Fossil Taxa

Luciano Varela^{1,2*}, P. Sebastián Tambusso^{1,2}

¹Departamento de Paleontología, Facultad de Ciencias, Universidad de la República, Iguá 4225, 11400 Montevideo, Uruguay

²Servicio Académico Universitario y Centro de Estudio Paleontológicos (SAUCE-P), Universidad de la República, Santa Isabel s/n, 91500 Sauce, Departamento de Canelones, Uruguay

*Corresponding author: luciano.lvr@gmail.com

Abstract

The present 3D Dataset contains the 3D models analyzed in 3D Finite Element Analysis and Geometric Morphometrics of Sloths (Xenarthra, Folivora) Mandibles Show Insights on the Dietary Specializations of Fossil Taxa. Journal of South American Earth Sciences. <https://doi.org/10.1016/j.jsames.2023.104445>.

Keywords: Ground Sloths, Mandibles, Photogrammetry, Quaternary, South America

Submitted:01/06/2023, published online:10/06/2023. <https://doi.org/10.18563/journal.m3.199>

INTRODUCTION

We present the surface models (Fig. 1; Table 1) of several specimens of Folivora (Mammalia, Xenarthra) used in Varela et al. (2023). The scanned specimens represent mandibles of adult individuals of representatives of one of the mayor clades of sloths, the Mylodontoidea (Delsuc et al. 2019; Varela et al. 2019), namely *Lestodon armatus*, *Glossotherium robustum*, *Myodon darwini*, *Scelidotherium leptcephalum*, and *Valgipes bucklandi*. The specimens also correspond the some of the most representative taxa of the Río de la Plata region during the Late Pleistocene (Varela and Fariña 2016; Varela et al. 2018). The models were used in Varela et al. (2023) to perform Finite Elements Analysis and Geometric Morphometrics to explore potential differences among taxa regarding their dietary specializations. The results supported the existence of adaptations to hard-food processing in some taxa and consequently the probable resource partitioning among members of the Mylodontoidea coexisting in the same region during the Late Pleistocene.

METHODS

The fossil specimens were scanned using photogrammetry with the software Agisoft Photoscan. The 3D surface models are provided in .stl format, which can be opened by an extensive list of free and open-source software.

ACKNOWLEDGEMENTS

We thank the curators of the following collections, who kindly allowed us access to the specimens under their care: Kasper Lykke Hansen (NHMD), Andrés Rinderknecht (MNHN-M); and the personnel of the museums of Colonia del Sacramento, Uruguay. This work was supported by a grant from the Espacio Interdisciplinario, Udelar (Núcleo interdisciplinario de estudios cuaternarios 2019) to RAF, and a grant by the Comisión Sectorial de Posgrado, Udelar and a Sepkoski Grant from the

Inv nr.	Taxon	Description	Collection
CAV 379	<i>Myodon darwini</i>	Right hemi-mandible	SAUCE-P, Uruguay
MNHN-M 137,722	<i>Scelidotherium leptcephalum</i>	Mandible	MNHN, Montevideo, Uruguay
MNHN-M 914	<i>Glossotherium robustum</i>	Mandible	MNHN, Montevideo, Uruguay
MPAC 899	<i>Lestodon armatus</i>	Mandible	MPAC, Colonia, Uruguay
NHMD.Z.M.K 1/1845:3540	<i>Valgipes bucklandi</i>	Mandible	NHMD, Copenhagen, Denmark

Table 1. Involved specimens. Collections: Statens Naturhistoriske Museum, Copenhagen, Denmark (NHMD); Museo Nacional de Historia Natural, Montevideo, Uruguay (MNHN-M); Museo Paleontológico Profesor Armando Calcaterra, Colonia, Uruguay (MPAC); Servicio Académico Universitario y Centro de Estudio Paleontológicos, Universidad de la República, Canelones, Uruguay (SAUCE-P).

Paleontological Society International Research Program (Pal-SIRP) to LV.

BIBLIOGRAPHY

- Delsuc, F., Kuch, M., Gibb, G. C., Karpinski, E., Hackenberger, D., Szpak, P., ... & Poinar, H. N. 2019. Ancient mitogenomes reveal the evolutionary history and biogeography of sloths. *Current Biology*, 29(12), 2031-2042. <https://doi.org/10.1016/j.cub.2019.05.043>
- Lobato, C., Varela, L., Tambusso, P. S., Miño-Boilini, Á. R., Clavijo, L., Fariña, R. A. 2021. Presence of the ground sloth *Valgipes bucklandi* (Xenarthra, Folivora, Scelidotheriinae) in southern Uruguay during the Late Pleistocene: Ecological and




Figure 1. 3D surface models of: CAV 379 (*Mylodon darwini*), MPAC 899 (*Lestodon armatus*), MNHN-M 914 (*Glossotherium robustum*), MNHN-M 137,722 (*Scelidotherium leptcephalum*), and NHMD.Z.M.K. 1/1845:3540 (*Valgipes bucklandi*) in lateral view. Scale bar: 10 cm.

- biogeographical implications. *Quaternary International*, 601, 104-115. <https://doi.org/10.1016/j.quaint.2021.06.011>
- Varela, L., Fariña, R. A., 2016. Co-occurrence of mylodontid sloths and insights on their potential distributions during the late Pleistocene. *Quat. Res.* 85 (1), 66-74. <https://doi.org/10.1016/j.yqres.2015.11.009>.
- Varela, L., Tambusso, P. S., Patiño, S. J., Di Giacomo, M., Fariña, R. A. 2018. Potential distribution of fossil xenarthrans in South America during the Late Pleistocene: Co-Occurrence and provincialism. *Journal of Mammalian Evolution*, 1-12. <https://doi.org/10.1007/s10914-017-9406-9>
- Varela, L., Tambusso, P. S., McDonald, H. G., Fariña, R. A. 2019. Phylogeny, macroevolutionary trends and historical biogeography of sloths: insights from a Bayesian morphological clock analysis. *Systematic Biology*, 68(2), 204-218. <https://doi.org/10.1093/sysbio/syy058>
- Varela, L., Tambusso, P. S., Pérez-Zerpa J. M., McAfee, R. K., Fariña, R. A. 2023. 3D Finite Element Analysis and Geometric Morphometrics of Sloths (Xenarthra, Folivora) Mandibles Show Insights on the Dietary Specializations of Fossil Taxa. *Journal of South American Earth Sciences*. <https://doi.org/10.1016/j.jsames.2023.104445>